

CLUSTER STRUCTURES FOR 2-CALABI-YAU CATEGORIES AND UNIPOTENT GROUPS

A. B. BUAN, O. IYAMA, I. REITEN, AND J. SCOTT

ABSTRACT. We investigate cluster tilting objects (and subcategories) in triangulated 2-Calabi-Yau categories and related categories. In particular we construct a new class of such categories related to preprojective algebras of non-Dynkin quivers associated with elements in the Coxeter group. This class of 2-Calabi-Yau categories contains the cluster categories and the stable categories of preprojective algebras of Dynkin graphs as special cases. For these 2-Calabi-Yau categories we construct cluster tilting objects associated with each reduced expression. The associated quiver is described in terms of the reduced expression. Motivated by the theory of cluster algebras, we formulate the notions of (weak) cluster structure and substructure, and give several illustrations of these concepts. We give applications to cluster algebras and subcluster algebras related to unipotent groups, both in the Dynkin and non-Dynkin case.

CONTENTS

Introduction	1
I. 2-CY categories and substructures	3
I.1. Cluster structures	3
I.2. Substructures	11
I.3. Preprojective algebras of Dynkin type	16
II. Preprojective algebras for non-Dynkin quivers	20
II.1. Tilting modules over 2-CY algebras	20
II.2. Cluster tilting objects for preprojective algebras	25
II.3. Realization of cluster categories and stable categories for preprojective algebras of Dynkin type	29
II.4. Quivers of cluster tilting subcategories	31
II.5. Substructure	36
III. Connections to cluster algebras	37
III.1. Cluster algebras, subcluster algebras and cluster maps	37
III.2. The GLS φ -map with applications to the Dynkin case	39
III.3. Cluster Structure of the loop group $SL_2(\mathcal{L})$	43
References	47

INTRODUCTION

The theory of cluster algebras, initiated by Fomin-Zelevinsky in [FZ2], and further developed in a series of papers, including [FZ3, BFZ, FZ4], has turned out to have interesting connections with many parts of algebra and other branches of mathematics. One of the links is with the representation theory of algebras, where a first connection was discovered in [MRZ]. A philosophy has been to model the main ingredients in the definition of a cluster algebra in a categorical/module theoretical setting. The cluster categories associated with

finite dimensional hereditary algebras were introduced for this purpose in [BMRRT], and shown to be triangulated in [Ke1] (see also [CCS] for the A_n case), and the module categories $\text{mod } \Lambda$ for Λ a preprojective algebra of a Dynkin quiver have been used for a similar purpose [GLS1]. This development has both inspired new directions of investigations on the categorical side, as well as interesting feedback on the theory of cluster algebras, see for example [ABS, BMR1, BMR2, BM, BMRT, CC, CK1, CK2, GLS1, GLS5, Hu, I1, I2, IR, IY, IT, KR1, KR2, Rin2, T] for material related to this paper.

Both the cluster categories and the stable categories $\underline{\text{mod}} \Lambda$ of preprojective algebras are triangulated Calabi-Yau categories of dimension 2 (2-CY for short). They both have what is called cluster tilting objects/subcategories [BMRRT, KR1, IY] (called maximal 1-orthogonal in [I1]), which are important since they are the analogs of clusters. The investigation of cluster tilting objects/subcategories in 2-CY categories and related categories is interesting both from the point of view of cluster algebras and in itself. Hence it is of interest to develop methods for constructing 2-CY categories together with the special objects/subcategories, and this is the main purpose of the first two chapters.

The properties of cluster tilting objects in (Hom-finite) 2-CY categories which have been important for applications to cluster algebras are (a) the unique exchange property for indecomposable summands of cluster tilting objects, (b) the existence of associated exchange triangles, (c) having no loops or 2-cycles (in the quiver of the endomorphism algebra of a cluster tilting object) and (d) when passing from the endomorphism algebra of a cluster tilting object T to the endomorphism algebra of another one T^* via an exchange, the change in quivers is given by Fomin-Zelevinsky mutation. The properties (a) and (b) are known to hold for any 2-CY triangulated category [IY], proved for cluster categories in [BMRRT] and for stable categories of preprojective algebras of Dynkin type in [GLS1]. The property (c) does not always hold (see [BIKR]), and hence it is of interest to establish criteria for this to be the case, which is one of the topics in this paper. We then show for any 2-CY categories that if (c) holds, then also (d) follows, as previously shown by Palu for algebraic triangulated categories [P]. We construct new 2-CY categories with cluster tilting objects from old ones via some subfactor construction, extending results from [IY], with a main focus on how condition (c) behaves under this construction. Associated with this we introduce the notions of cluster structures and cluster substructures.

Important examples, investigated in [GLS1], are the categories $\text{mod } \Lambda$ of finitely generated modules over the preprojective algebra Λ of a Dynkin quiver. We deal with appropriate subcategories of $\text{mod } \Lambda$. The main focus in this paper is on the more general case of subcategories of the category $\text{f.l. } \Lambda$ of finite length modules over the completion of the preprojective algebra of a non-Dynkin quiver with no loops. Our main tool is to extend the tilting theory developed for Λ in the noetherian case in [IR]. This turns out to give a large class of 2-CY categories associated with elements in the corresponding Coxeter groups. For these categories we construct cluster tilting objects associated with each reduced expression, and we describe the associated quiver directly in terms of the reduced expression. We prove that this class of 2-CY categories contains all the cluster categories of finite dimensional hereditary algebras and the stable categories $\underline{\text{mod}} \Lambda$ for a preprojective algebra Λ of Dynkin type. This also allows us to get more information on the latter case.

We illustrate with applications to constructing subcluster algebras of cluster algebras, a notion which we define here, and which is already implicit in the literature. For this we define, inspired by maps from [CC, CK1] and [GLS1], (strong) cluster maps. These maps have the property that we can pass from cluster structures and substructures to cluster algebras and subcluster algebras.

Associated with substructures for preprojective algebras of Dynkin type, we discuss examples from $\text{SO}_8(\mathbb{C})$ -isotropic Grassmanians, for the $G_{2,5}$ Schubert variety and for a unipotent cell of the unipotent subgroup of $\text{SL}_4(\mathbb{C})$. For preprojective algebras of extended Dynkin

type we use our results to investigate cluster structures for affine unipotent cells, in some special cases for the loop group $SL_2(\mathcal{L})$.

For a (non-Dynkin) quiver Q with associated Coxeter group W we can for each $w \in W$ consider the coordinate ring $\mathbb{C}[U^w]$ of the unipotent cell associated with w in the corresponding Kac-Moody group. We conjecture that this ring has a cluster algebra structure, and that it is modelled by our (stably) 2-CY category associated with the same w . As a support for this we prove the conjecture for the case $\widehat{A}_1$ for the word w of length at most 4.

The first chapter is devoted to introducing and investigating the notions of cluster structures and substructures, and giving sufficient conditions for such structures to occur. Also the three concrete examples mentioned above are investigated, and used to illustrate the connection with cluster algebras and subcluster algebras in Section III.2. In Chapter II we use tilting theory to construct categories whose stable categories are 2-CY, along with natural cluster tilting objects in these categories. In Section III.2 we illustrate with examples for preprojective algebras of Dynkin type, and in Section III.3 we illustrate with examples from the extended Dynkin case.

Part of this work was done independently by Geiss-Leclerc-Schröer in [GLS5]. For Chapter I, this concerns the development of 2-CY categories (in a different language) in the case of subcategories of the form $\text{Sub } P$ (or $\text{Fac } P$) for P projective, over a preprojective algebra of Dynkin type, with a somewhat different approach. Concerning Section III.2, examples arising from $\text{Sub } P$ were done independently in [GLS5]. For this connection, the last author was inspired by a lecture of Leclerc in 2005, where cluster algebras associated with $\text{Sub } P$ in the A_n -case were discussed. There is also recent work of Geiss-Leclerc-Schröer [GLS6] related to Chapter II, where completely different methods are used.

For general background on representation theory of algebras, we refer to [ARS, ASS, Rin1, H1, AHK], and for Lie theory we refer to [BL].

Our modules are usually left modules and composition of maps fg means first f , then g .

The second author would like to thank William Crawley-Boevey and Christof Geiss for answering a question on references about 2-CY property of preprojective algebras. He also would like to thank Bernald Leclerc for valuable comments.

I. 2-CY CATEGORIES AND SUBSTRUCTURES

The cluster algebras of Fomin and Zelevinsky have motivated work on trying to model the essential ingredients in the definition of a cluster algebra in a categorical/module theoretical way. In particular, this led to the theory of cluster categories and the investigation of new aspects of the module theory of preprojective algebras of Dynkin type. In Section I.1 we give some of the main categorical requirements needed for the modelling, for the cases with and without coefficients, leading to the notions of weak cluster structure and cluster structure. The main examples do, like the above mentioned examples, have 2-Calabi-Yau type properties.

We introduce substructures of (weak) cluster structures in Section I.2. For this it is natural to deal with (weak) cluster structures with what will be called coefficients, at least for the substructures. Of particular interest for our applications to cluster algebras is the case of completions of preprojective algebras Λ of a finite connected quiver with no loops over an algebraically closed field k , where the interesting larger category is the stable category $\underline{\text{f.l.}}\Lambda$ of the finite length Λ -modules. For Dynkin quivers this is the stable category $\underline{\text{mod}}\Lambda$ of the finitely generated Λ -modules, and in the non-Dynkin case $\underline{\text{f.l.}}\Lambda = \text{f.l.}\Lambda$. The first case is discussed in Section I.3, while Chapter II is devoted to the second case.

I.1. Cluster structures.

In this section we introduce the concepts of weak cluster structure and cluster structure for extension closed subcategories of triangulated categories or for exact categories. We

illustrate with 2-CY categories and closely related categories, and the main objects we investigate are the cluster tilting ones. These cases are particularly nice when the quivers of the cluster tilting subcategories have no loops or 2-cycles. Also the closely related maximal rigid objects (see [GLS1]) provide interesting examples.

We start with introducing the notions of weak cluster structure and cluster structure. Throughout this chapter all categories are Krull-Schmidt categories over an algebraically closed field k , that is, each object is isomorphic to a finite direct sum of indecomposable objects with local endomorphism ring. The categories we consider are either exact (for example abelian) categories or extension closed subcategories of triangulated categories. Note that an extension closed subcategory of an exact category is again exact. We refer to [Ke2, Ke3] for definition and basic properties of exact categories, which behave very much like abelian categories, also with respect to derived categories and Ext-functors.

We often identify a set of indecomposable objects with the additive subcategory consisting of all summands of direct sums of these indecomposable objects. We also identify an object with the set of indecomposable objects appearing in a direct sum decomposition, and with the subcategory obtained in the above way.

Assume that we have a collection of sets $\underline{x}$ (which may be infinite) of non-isomorphic indecomposable objects, called *clusters*. The union of all indecomposable objects in clusters are called *cluster variables*. Assume also that there is a subset $\underline{p}$ (which may be infinite) of indecomposable objects which are not cluster variables, called *coefficients*. We denote by T the union of the indecomposable objects in $\underline{x}$ and $\underline{p}$, sometimes viewed as a category with these objects, and call it an *extended cluster*.

We say that the clusters, together with the prescribed set of coefficients $\underline{p}$, give a *weak cluster structure* on $\mathcal{C}$ if the following hold:

- (a) For each extended cluster T and each cluster variable M which is a summand in T , there is a unique indecomposable object $M^* \not\cong M$ such that we get a new extended cluster T^* by replacing M by M^* . We denote this operation, called *exchange*, by $\mu_M(T) = T^*$, and we call (M, M^*) an *exchange pair*.
- (b) There are triangles/short exact sequences $M^* \xrightarrow{f} B \xrightarrow{g} M$ and $M \xrightarrow{s} B' \xrightarrow{t} M^*$, where the maps g and t are minimal right $\text{add}(T \setminus \{M\})$ -approximations and f and s are minimal left $\text{add}(T \setminus \{M\})$ -approximations. These are called *exchange triangles/sequences*.

Denote by Q_T the quiver of T , where the vertices correspond to the indecomposable objects in T and the number of arrows $T_i \rightarrow T_j$ between two indecomposable objects T_i and T_j is given by the dimension of the space of irreducible maps $\text{rad}(T_i, T_j) / \text{rad}^2(T_i, T_j)$. Here $\text{rad}(\ , \)$ denotes the radical in $\text{add } T$, where the objects are finite direct sums of objects in T . For an algebra Λ (where Λ has a unique decomposition as a direct sum of indecomposable objects), the quiver of Λ is then the opposite of the quiver of $\text{add } \Lambda$.

We say that a quiver $Q = (Q_0, Q_1)$ is an *extended quiver* with respect to a subset of vertices Q'_0 if there are no arrows between two vertices in $Q_0 \setminus Q'_0$. We regard the quiver Q_T of an extended cluster as an extended quiver by neglecting all arrows between two vertices corresponding to coefficients.

We say that $\mathcal{C}$, with a fixed set of clusters and coefficients, has *no loops* (respectively, *no 2-cycles*) if in the extended quiver of each extended cluster there are no loops (respectively, no 2-cycles). When $\underline{x}$ is finite, this is the opposite quiver of the factor algebra $\underline{\text{End}}(T)$ of $\text{End}(T)$ by the maps factoring through direct sums of objects from $\underline{p}$.

We say that we have a *cluster structure* if the following additional conditions hold:

- (c) There are no loops or 2-cycles. (In other words, for a cluster variable M , any non-isomorphism $u: M \rightarrow M$ factors through $g: B \rightarrow M$ and through $s: M \rightarrow B'$, and any non-isomorphism $v: M^* \rightarrow M^*$ factors through $f: M^* \rightarrow B$ and through $t: B' \rightarrow M^*$, and B and B' have no common indecomposable summand.)

- (d) For an extended cluster T , passing from Q_T to Q_{T^*} is given by Fomin-Zelevinsky mutation at the vertex of Q_T given by the cluster variable M .

Note that (c) is needed for (d) to make sense, but it is still convenient to write two separate statements.

We recall that for an extended quiver Q without loops or 2-cycles and a vertex i in Q'_0 , the Fomin-Zelevinsky mutation $\mu_i(Q)$ of Q at i is the quiver obtained from Q making the following changes [FZ2]:

- Reverse all the arrows starting or ending at i .
- Let $s \neq i$ and $t \neq i$ be vertices in Q_0 such that at least one vertex belongs to Q'_0 . If we have $n > 0$ arrows from t to i and $m > 0$ arrows from i to s in Q and r arrows from s to t in Q (interpreted as $-r$ arrows from t to s if $r < 0$), then we have $nm - r$ arrows from t to s in the new quiver $\mu_i(Q)$ (interpreted as $r - nm$ arrows from s to t if $nm - r < 0$).

The main known examples of triangulated k -categories with finite dimensional homomorphism spaces (Hom-finite for short) which have a weak cluster structure, and usually cluster structure, are 2-CY categories. These are triangulated k -categories with functorial isomorphisms $D \operatorname{Ext}^1(A, B) \simeq \operatorname{Ext}^1(B, A)$ for all A, B in $\mathcal{C}$, where $D = \operatorname{Hom}_k(-, k)$. A Hom-finite triangulated category is 2-CY if and only if it has almost split triangles with translation τ and $\tau: \mathcal{C} \rightarrow \mathcal{C}$ is a functor isomorphic to the shift functor $[1]$ (see [RV]).

We have the following examples of 2-CY categories.

- (1) The cluster category $\mathcal{C}_H$ associated with a finite dimensional hereditary k -algebra H is by definition the orbit category $\mathbf{D}^b(H)/\tau^{-1}[1]$, where $\mathbf{D}^b(H)$ is the bounded derived category of finitely generated H -modules, and τ is the AR-translation of $\mathbf{D}^b(H)$ [BMRR]. It is a Hom-finite triangulated category [Ke2], and it is 2-CY since $\tau = [1]$.

- (2) The stable category of maximal Cohen-Macaulay modules $\underline{\mathbf{CM}}(R)$ over a 3-dimensional complete local commutative noetherian Gorenstein isolated singularity R containing the residue field k [A2] (see [Yo]).

- (3) The preprojective algebra Λ associated to a finite connected quiver Q without loops is defined as follows: Let $\tilde{Q}$ be the quiver constructed from Q by adding an arrow $\alpha^*: i \rightarrow j$ for each arrow $\alpha: j \rightarrow i$ in Q . Then $\Lambda = k\tilde{Q}/I$, where I is the ideal generated by the sum of commutators $\sum_{\beta \in Q_1} [\beta, \beta^*]$. Note that Λ is uniquely determined up to isomorphism by the underlying graph of Q .

When Λ is the preprojective algebra of a Dynkin quiver over k , the stable category $\underline{\mathbf{mod}}\Lambda$ is 2-CY (see [AR2, 3.1.1.2][CB][Ke2, 8.5]).

When Λ is the completion of the preprojective algebra of a finite connected quiver without loops which is not Dynkin, the bounded derived category $\mathbf{D}^b(\text{f.l. } \Lambda)$ of the category f.l. Λ of the modules of finite length is 2-CY (see [B, CB, BBK][GLS4, section 8]).

We shall also use the terminology 2-CY in more general situations. Note that from now on we will usually write just “category” instead of “ k -category”.

We say that an exact Hom-finite category $\mathcal{C}$ is *derived 2-CY* if the triangulated category $\mathbf{D}^b(\mathcal{C})$ is 2-CY, i.e. if $D \operatorname{Ext}^i(A, B) \simeq \operatorname{Ext}^{2-i}(B, A)$ for all A, B in $\mathbf{D}^b(\mathcal{C})$ and all i . Note that when $\mathcal{C}$ is derived 2-CY, then $\mathcal{C}$ has no non-zero projective or injective objects. The category f.l. Λ where Λ is the completion of the preprojective algebra of a non-Dynkin connected quiver without loops is an important example of a derived 2-CY category.

We say that an exact category $\mathcal{C}$ is *stably 2-CY* if it is Frobenius, that is, $\mathcal{C}$ has enough projectives and injectives, which coincide, and the stable category $\underline{\mathcal{C}}$, which is triangulated [H1], is Hom-finite 2-CY. Recall that $\mathcal{C}$ is said to have enough projectives if for each X in $\mathcal{C}$ there is an exact sequence $0 \rightarrow Y \rightarrow P \rightarrow X \rightarrow 0$ in $\mathcal{C}$ with P projective. Having enough injectives is defined in a dual way.

We have the following characterization of stably 2-CY categories.

Proposition I.1.1. *Let $\mathcal{C}$ be an exact Frobenius category. Then $\mathcal{C}$ is stably 2-CY if and only if $\text{Ext}_{\mathcal{C}}^1(A, B)$ is finite dimensional and we have functorial isomorphisms $D\text{Ext}_{\mathcal{C}}^1(A, B) \simeq \text{Ext}_{\mathcal{C}}^1(B, A)$ for all A, B in $\mathcal{C}$.*

Proof. Let A and B be in $\mathcal{C}$, and let $0 \rightarrow A \rightarrow P \rightarrow \Omega^{-1}A \rightarrow 0$ be an exact sequence in $\mathcal{C}$ where P is projective injective. Apply $\text{Hom}_{\mathcal{C}}(B, _)$ to get the exact sequence $0 \rightarrow \text{Hom}_{\mathcal{C}}(B, A) \rightarrow \text{Hom}_{\mathcal{C}}(B, P) \rightarrow \text{Hom}_{\mathcal{C}}(B, \Omega^{-1}A) \rightarrow \text{Ext}_{\mathcal{C}}^1(B, A) \rightarrow 0$. Then we get $\text{Ext}_{\mathcal{C}}^1(B, A) \simeq \text{Hom}_{\mathcal{C}}(B, \Omega^{-1}A) = \text{Ext}_{\mathcal{C}}^1(B, A)$.

Assume that $\mathcal{C}$ is stably 2-CY, that is, the stable category $\underline{\mathcal{C}}$ is a Hom-finite triangulated 2-CY category. Then $\text{Ext}_{\underline{\mathcal{C}}}^1(B, A)$ is finite dimensional for all A, B in $\mathcal{C}$, and hence $\text{Ext}_{\mathcal{C}}^1(B, A)$ is finite dimensional, and we have functorial isomorphisms $D\text{Ext}_{\mathcal{C}}^1(A, B) \simeq \text{Ext}_{\mathcal{C}}^1(B, A)$.

The converse also follows directly. $\square$

Examples of exact stably 2-CY categories are categories of maximal Cohen-Macaulay modules $\text{CM}(R)$ for a 3-dimensional complete local commutative isolated Gorenstein singularity R (containing the residue field k) and $\text{mod } \Lambda$ for Λ being the preprojective algebra of a Dynkin quiver. We shall see several further examples later.

We are especially interested in pairs of 2-CY categories $(\mathcal{C}, \underline{\mathcal{C}})$ where $\mathcal{C}$ is an exact stably 2-CY category. The only difference in indecomposable objects between $\mathcal{C}$ and $\underline{\mathcal{C}}$ is the indecomposable projective objects in $\mathcal{C}$. Also note that given an exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ in $\mathcal{C}$, there is an associated triangle $A \rightarrow B \rightarrow C \rightarrow A[1]$ in $\underline{\mathcal{C}}$. Conversely, given a triangle $A \rightarrow B \xrightarrow{g} C \rightarrow A[1]$ in $\underline{\mathcal{C}}$, we lift $\underline{g} \in \text{Hom}_{\underline{\mathcal{C}}}(B, C)$ to $g \in \text{Hom}_{\mathcal{C}}(B, C)$, and obtain an exact sequence $0 \rightarrow A \rightarrow B \oplus P \rightarrow C \rightarrow 0$ in $\mathcal{C}$, where P is projective. We then have the following useful fact.

Proposition I.1.2. *Let $\mathcal{C}$ be an exact stably 2-CY category with a set of clusters $\underline{x}$ and a set of coefficients $\underline{p}$ which are the indecomposable projective objects. For the stable 2-CY category $\underline{\mathcal{C}}$ consider the same set of clusters $\underline{x}$ and with no coefficients. Then we have the following.*

- (a) *The $(\underline{x}, \underline{p})$ give a weak cluster structure on $\mathcal{C}$ if and only if the $(\underline{x}, \emptyset)$ give a weak cluster structure on $\underline{\mathcal{C}}$.*
- (b) *$\mathcal{C}$ has no loops if and only if $\underline{\mathcal{C}}$ has no loops.*
- (c) *If $\mathcal{C}$ has a cluster structure, then $\underline{\mathcal{C}}$ has a cluster structure.*

Proof. (a) Assume we have a collection of extended clusters for $\mathcal{C}$. Assume also that $\mu_M(T)$ is defined for each indecomposable non-projective object M in the cluster T , and that we have the required exchange exact sequences. Then the induced clusters for $\underline{\mathcal{C}}$ determine a weak cluster structure for $\underline{\mathcal{C}}$. The converse also follows directly.

(b) For an extended cluster T in $\mathcal{C}$, the quiver $\bar{Q}_T$ is obtained from the quiver Q_T by removing the vertices corresponding to the indecomposable projective objects. The claim is then obvious.

(c) It is clear that if there is no 2-cycle for $\mathcal{C}$, then there is no 2-cycle for $\underline{\mathcal{C}}$, and the claim follows from this. $\square$

In the examples of 2-CY categories with cluster structure which have been investigated the extended clusters have been the subcategories T where $\text{Ext}^1(M, M) = 0$ for all $M \in T$, and whenever $X \in \mathcal{C}$ satisfies $\text{Ext}^1(M, X) = 0$ for all $M \in T$, then $X \in T$. Such T has been called *cluster tilting subcategory* in [BMRRT, KR1] if it is in addition functorially finite in the sense of [AS], which is automatically true when T is finite. Such T has been called maximal 1-orthogonal subcategory in [I1, I2], and Ext-configuration in [BMRRT], without the assumption on functorially finiteness.

We have the following nice connections between $\mathcal{C}$ and $\underline{\mathcal{C}}$ for an exact stably 2-CY category $\mathcal{C}$ when using the cluster tilting subcategories.

Lemma I.1.3. *Let $\mathcal{C}$ be an exact stably 2-CY category, and let T be a subcategory of $\mathcal{C}$ containing all indecomposable projective objects. Then T is a cluster tilting subcategory in $\mathcal{C}$ if and only if it is the same in $\underline{\mathcal{C}}$.*

Proof. We have $\text{Ext}_{\mathcal{C}}^1(C, A) \simeq \text{Ext}_{\underline{\mathcal{C}}}^1(C, A)$ from the proof of Proposition I.1.2. It is easy to see that T is functorially finite in $\mathcal{C}$ if and only if it is functorially finite in $\underline{\mathcal{C}}$ [AS]. Hence T is cluster tilting in $\mathcal{C}$ if and only if it is cluster tilting in $\underline{\mathcal{C}}$. $\square$

Lemma I.1.4. *Let $\mathcal{C}$ be an exact stably 2-CY category and T a cluster tilting object in $\mathcal{C}$, with an indecomposable non-projective summand M . Then there is no loop at M for $\text{End}_{\mathcal{C}}(T)$ if and only if there is no loop at M for $\text{End}_{\underline{\mathcal{C}}}(T)$. If $\mathcal{C}$ has no 2-cycles, there are none for $\underline{\mathcal{C}}$.*

Proof. This is a direct consequence of Proposition I.1.2 and the definitions. $\square$

Note that an exact stably 2-CY category $\mathcal{C}$, with the cluster tilting subcategories, gives a situation where we have a natural set of coefficients, namely the indecomposable projective objects which clearly belong to all cluster tilting subcategories, whereas $\underline{\mathcal{C}}$ with the cluster tilting subcategories gives a case where it is natural to choose no coefficients. We have the following useful observation, which follows from Proposition I.1.2.

Proposition I.1.5. *Let $\mathcal{C}$ be a Hom-finite exact stably 2-CY category. Then the cluster tilting subcategories in $\mathcal{C}$, with the indecomposable projectives as coefficients, determine a weak cluster structure on $\mathcal{C}$ if and only if the cluster tilting subcategories in $\underline{\mathcal{C}}$ determine a weak cluster structure on $\underline{\mathcal{C}}$.*

When $\mathcal{C}$ is Hom-finite triangulated 2-CY, then $\mathcal{C}$ has a weak cluster structure, with the extended clusters being the cluster tilting subcategories and the indecomposable projectives being the coefficients [IY]. Properties (c) and (d) hold for cluster categories and the stable category $\underline{\text{mod}}\Lambda$ of a preprojective algebra of Dynkin type [BMRRT, BMR2, GLS1], but (c) does not hold in general [BIKR]. However, we show that when we have some cluster tilting object in the 2-CY category $\mathcal{C}$, then (d) holds under the assumption that (c) holds. This was first proved in [P] when $\mathcal{C}$ is *algebraic*, that is by definition the stable category of a Frobenius category, as a special case of a more general result. Our proof is inspired by [IR, 7.1].

Theorem I.1.6. *Let $\mathcal{C}$ be Hom-finite triangulated (or exact stably) 2-CY category with some cluster tilting subcategory. If $\mathcal{C}$ has no loops or 2-cycles, then the cluster tilting subcategories determine a cluster structure for $\mathcal{C}$.*

Proof. We give a proof for the triangulated 2-CY case. Using exact sequences instead of triangles, a similar argument works for the stably 2-CY case. Note that in the stably 2-CY case we do not have to consider arrows between projective vertices.

Let $T = \bigoplus_{i=1}^n T_i$ be a cluster tilting subcategory in $\mathcal{C}$. Fix a vertex $k \in \{1, \dots, n\}$, and let $T^* = \bigoplus_{i \neq k} T_i \oplus T_k^* = \mu_k(T)$. We have exchange triangles $T_k^* \rightarrow B_k \rightarrow T_k$ and $T_k \rightarrow B'_k \rightarrow T_k^*$, showing that when passing from $\text{End}(T)$ to $\text{End}(T^*)$ we reverse all arrows in the quiver of $\text{End}(T)$ starting or ending at k .

We need to consider the situation where we have arrows $j \rightarrow k \rightarrow i$. Since there are no 2-cycles, there is no arrow $i \rightarrow k$. Consider the exchange triangles $T_i^* \rightarrow B_i \rightarrow T_i$ and $T_i \rightarrow B'_i \rightarrow T_i^*$. Then T_k is not a direct summand of B'_i , and we write $B_i = D_i \oplus T_k^m$ for some $m > 0$, where T_k is not a direct summand of D_i .

Starting with the maps in the upper square and the triangles they induce, we get by the octahedral axiom the diagram below, where the third row is a triangle.

$$\begin{array}{ccccccc}
 & & (T_k^*)^m[1] & \xlongequal{\quad} & (T_k^*)^m[1] & & \\
 & & \uparrow & & \uparrow & & \\
 T_i[-1] & \longrightarrow & T_i^* & \longrightarrow & D_i \oplus T_k^m & \longrightarrow & T_i \\
 \parallel & & \uparrow & & \uparrow & & \parallel \\
 T_i[-1] & \longrightarrow & X & \longrightarrow & D_i \oplus B_k^m & \longrightarrow & T_i \\
 & & \uparrow & & \uparrow & & \\
 & & (T_k^*)^m & \xlongequal{\quad} & (T_k^*)^m & &
 \end{array}$$

Using again the octahedral axiom, we get the following commutative diagram of triangles, where the second row is an exchange triangle and the third column is the second column of the previous diagram.

$$\begin{array}{ccccccc}
 & & (T_k^*)^m[1] & \xlongequal{\quad} & (T_k^*)^m[1] & & \\
 & & \uparrow & & \uparrow & & \\
 T_i & \longrightarrow & B'_i & \longrightarrow & T_i^* & \longrightarrow & T_i[1] \\
 \parallel & & \uparrow & & \uparrow & & \parallel \\
 T_i & \longrightarrow & Y & \longrightarrow & X & \longrightarrow & T_i[1] \\
 & & \uparrow & & \uparrow & & \\
 & & (T_k^*)^m & \xlongequal{\quad} & (T_k^*)^m & &
 \end{array}$$

Since T_k is not in $\text{add } B'_i$, we have $(B'_i, (T_k^*)^m[1]) = 0$, and hence $Y = B'_i \oplus (T_k^*)^m$.

Consider the triangle $X \rightarrow D_i \oplus B_k^m \xrightarrow{a} T_i \rightarrow X[1]$. Let $\overline{T}^* = (\oplus_{t \neq i, k} T_t) \oplus T_k^*$. We observe that $D_i \oplus B_k^m$ is in $\text{add } \overline{T}^*$. For $D_i \oplus B_k^m = B_i$ is in $\text{add } T$. Since there is no loop at i , then T_i is not a direct summand of D_i , and T_i is not a direct summand of B_k since there is no arrow from i to k . Further T_k is not a direct summand of D_i by the choice of D_i , and T_k is not a direct summand of B_k since there is no loop at k . Hence we see that B_i is in $\text{add } \overline{T}^*$.

We next want to show that a is a right $\text{add } \overline{T}^*$ -approximation. It follows from the first commutative diagram that any map $g: T_t \rightarrow T_i$, where T_t is an indecomposable direct summand of $\overline{T}^*$ not isomorphic to T_k^* , factors through a . Let then $f: T_k^* \rightarrow T_i$ be a map, and $h: T_k^* \rightarrow B_k$ the minimal left $\text{add } \overline{T}$ -approximation, where $\overline{T} = \oplus_{t \neq k} T_t$. Then there is some $s: B_k \rightarrow T_i$ such that $hs = f$. Then s factors through a by the above, since B_k is in $\text{add } \overline{T}^*$ (using that T_i is not a direct summand in B_k), and T_k^* is not a direct summand of B_k . It follows that a is a right $\text{add } \overline{T}^*$ -approximation.

Consider now the triangle $T_i \rightarrow B'_i \oplus (T_k^*)^m \xrightarrow{b} X \rightarrow T_i[1]$. Then $B'_i \oplus (T_k^*)^m$ is clearly in $\text{add } \overline{T}^*$, since T_k is not a direct summand of B'_i . Since T_i is in both T and T^* , we have that $\text{Hom}(\overline{T}^*, T_i[1]) = 0$, and hence b is a right $\text{add } \overline{T}^*$ -approximation. Note that the approximations a and b need not be minimal.

Recall that we are interested in paths of length two $j \rightarrow k \rightarrow i$ passing through k . By the above, the number of arrows from j to i in the quiver Q_{T^*} is

$$u = \alpha_{D_i \oplus B_k^m}(T_j) - \alpha_{B'_i \oplus (T_k^*)^m}(T_j)$$

where $\alpha_X(T_j)$ denotes the multiplicity of T_j in X . We have


$$u = \alpha_{D_i}(T_j) + m\alpha_{B_k}(T_j) - \alpha_{B'_i}(T_j) = \alpha_{B_i}(T_j) + m\alpha_{B_k}(T_j) - \alpha_{B'_i}(T_j),$$

since $B_i = D_i \oplus T_k^m$. The last expression says that u is equal to the number of arrows from j to i in Q_T , minus the number of arrows from i to j , plus the product of the number of arrows from j to k and from k to i . This is what is required for having the Fomin-Zelevinsky mutation, and we are done. $\square$

We shall also use the terminology stably 2-CY for certain subcategories of triangulated categories. Let $\mathcal{B}$ be a functorially finite extension closed subcategory of a Hom-finite triangulated 2-CY category $\mathcal{C}$. We say that $X \in \mathcal{B}$ is *projective* in $\mathcal{B}$ if $\text{Hom}(X, \mathcal{B}[1]) = 0$. In this setting we shall prove in I.2.1 that the category $\mathcal{B}$ modulo projectives in $\mathcal{B}$ has a 2-CY triangulated structure. Note that $\mathcal{B}$ does not necessarily have enough projectives or injectives, for example if $\mathcal{B} = \mathcal{C}$. We then say that $\mathcal{B}$ is *stably 2-CY*.

We illustrate the concept with the following.

Example. Let $\mathcal{C}_Q$ be the cluster category of the path algebra kQ , where Q is the quiver $1 \xrightarrow{1} 2 \xrightarrow{3} 3$. We have the following AR-quiver for $\mathcal{C}_Q$, where S_i and P_i denote the simple and projective modules associated with vertex i respectively.



Then $\mathcal{B} = \text{mod } kQ$ is an extension closed subcategory of $\mathcal{C}_Q$ and it is easy to see that P_1 is the only indecomposable projective object in $\mathcal{B}$. Then $\mathcal{B}/P_1$ is clearly equivalent to the cluster category $\mathcal{C}_{Q'}$ where Q' is a quiver of type A_2 , which is a triangulated 2-CY category. Hence $\mathcal{B}$ is stably 2-CY.

In addition to the cluster tilting objects, also the maximal rigid objects have played an important role in the investigation of 2-CY categories. We now investigate the concepts of cluster structure and weak cluster structure with respect to these objects.

Recall that a subcategory T of a category $\mathcal{C}$ is said to be *rigid* if $\text{Ext}^1(M, M) = 0$ for all M in T , and *maximal rigid* if T is maximal among rigid subcategories [GLS1]. It is clear that any cluster tilting subcategory is maximal rigid, but the converse is not the case [BIKR]. There always exists a maximal rigid subcategory in $\mathcal{C}$ if the category $\mathcal{C}$ is skeletally small, while the existence of a cluster tilting subcategory is rather restrictive. It is of interest to have sufficient conditions for the two concepts to coincide. For this the following is useful (see [BMR1, I1, KR1] for (a) and the argument in [GLS1, 5.2] for (b)).

Proposition I.1.7. *Let $\mathcal{C}$ be a triangulated (or exact stably) 2-CY category.*

- (a) *Let T be a cluster tilting subcategory. Then for any X in $\mathcal{C}$, there exist triangles (or short exact sequences) $T_1 \rightarrow T_0 \rightarrow X$ and $X \rightarrow T'_0 \rightarrow T'_1$ with T_i, T'_i in T .*
- (b) *Let T be a functorially finite maximal rigid subcategory. Then for any X in $\mathcal{C}$ which is rigid, the same conclusion as in (a) holds.*

Then we have the following.

Theorem I.1.8. *Let $\mathcal{C}$ be an exact stably 2-CY category, with some cluster tilting object.*

- (a) *Then any maximal rigid object in $\mathcal{C}$ (respectively, $\underline{\mathcal{C}}$) is a cluster tilting object.*

- (b) Any rigid subcategory in $\mathcal{C}$ (respectively, $\underline{\mathcal{C}}$) has an additive generator which is a direct summand of a cluster tilting object.
- (c) All cluster tilting objects in $\mathcal{C}$ (respectively, $\underline{\mathcal{C}}$) have the same number of non-isomorphic indecomposable summands.

Proof. (a) Let N be maximal rigid in $\mathcal{C}$. We only have to show that any $X \in \mathcal{C}$ satisfying $\text{Ext}^1(N, X) = 0$ is contained in $\text{add } N$.

(i) Let M be a cluster tilting object in $\mathcal{C}$. Since N is maximal rigid and M is rigid, there exists an exact sequence $0 \rightarrow N_1 \rightarrow N_0 \rightarrow M \rightarrow 0$ with $N_i \in \text{add } N$ by Proposition I.1.7(b). In particular, we have $\text{pd}_{\text{End}(N)} \text{Hom}(N, M) \leq 1$.

(ii) Since M is cluster tilting, there is, by Proposition I.1.7(a), an exact sequence $0 \rightarrow X \rightarrow M_0 \rightarrow M_1 \rightarrow 0$ for X as above, with $M_i \in \text{add } M$, obtained by taking the minimal left $\text{add } M$ -approximation $X \rightarrow M_0$. Applying $(N, _)$, we have an exact sequence $0 \rightarrow (N, X) \rightarrow (N, M_0) \rightarrow (N, M_1) \rightarrow \text{Ext}^1(N, X) = 0$. By (i), we have $\text{pd}_{\text{End}(N)} \text{Hom}(N, X) \leq 1$. Take a projective resolution $0 \rightarrow (N, N_1) \rightarrow (N, N_0) \rightarrow (N, X) \rightarrow 0$. Then we have a complex

$$(1) \quad 0 \rightarrow N_1 \rightarrow N_0 \rightarrow X \rightarrow 0$$

in $\mathcal{C}$. Since $0 \rightarrow (P, N_1) \rightarrow (P, N_0) \rightarrow (P, X) \rightarrow 0$ is exact for any projective P in $\mathcal{C}$, it follows from the axioms of Frobenius categories that the complex (1) is an exact sequence in $\mathcal{C}$. Since $\text{Ext}^1(X, N) = 0$, we have $X \in \text{add } N$, and hence N is cluster tilting.

(b) Let M be a cluster tilting object in $\mathcal{C}$ and N a rigid object in $\mathcal{C}$. By [I2, 5.3.1], $\text{Hom}(M, N)$ is a partial tilting $\text{End}(M)$ -module. In particular, the number of non-isomorphic indecomposable direct summands of N is not greater than that of M . Consequently, any rigid object in $\mathcal{C}$ is a direct summand of some maximal rigid object in $\mathcal{C}$, which is cluster tilting by (a).

(c) See [I2, 5.3.3]. □

For a Hom-finite triangulated 2-CY category we also get a weak cluster structure, and sometimes a cluster structure, determined by the maximal rigid objects, if there are any. Note that there are cases where the maximal rigid objects are not cluster tilting [BIKR]. But we suspect the following.

Conjecture I.1.9. *Let $\mathcal{C}$ be a connected Hom-finite triangulated 2-CY category. Then any maximal rigid object without loops or 2-cycles in its quiver is a cluster tilting object.*

Furthermore, we have the following.

Theorem I.1.10. *Let $\mathcal{C}$ be a Hom-finite triangulated 2-CY category (or exact stably 2-CY category) having some functorially finite maximal rigid subcategory.*

- (a) *The functorially finite maximal rigid subcategories determine a weak cluster structure on $\mathcal{C}$.*
- (b) *If there are no loops or 2-cycles for the functorially finite maximal rigid subcategories, then they determine a cluster structure on $\mathcal{C}$.*

Proof. (a) This follows from [IY, 5.1, 5.3]. Note that the arguments there are stated only for cluster tilting subcategories, but work also for functorially finite maximal rigid subcategories.

(b) The proof of Theorem I.1.6 works also in this setting. □

There exist triangulated or exact categories with cluster tilting objects also when the categories are not 2-CY or stably 2-CY (see [I1, KZ, EH]). But we do not necessarily have even a weak cluster structure in this case. For let Λ be a Nakayama algebra with two simple modules S_1 and S_2 , with associated projective covers P_1 and P_2 . Assume first that P_1 and P_2 have length 3. Then in $\text{mod } \Lambda$ we have that $S_1 \oplus P_1 \oplus P_2$, $\frac{S_1}{S_2} \oplus P_1 \oplus P_2$, $\frac{S_2}{S_1} \oplus P_1 \oplus P_2$ are the cluster tilting objects, so we do not have the unique exchange property.

If P_1 and P_2 have length 4, then the cluster tilting objects are $S_1 \oplus \begin{smallmatrix} S_1 \\ S_2 \\ S_1 \end{smallmatrix} \oplus P_1 \oplus P_2$ and $S_2 \oplus \begin{smallmatrix} S_2 \\ S_1 \\ S_2 \end{smallmatrix} \oplus P_1 \oplus P_2$, and so there is no way of exchanging S_1 in the first object to obtain a new cluster tilting object.

We end this section with some information on the endomorphism algebras of cluster tilting objects in stably 2-CY categories. Such algebras are studied as analogs of Auslander algebras in [GLS1, I1, I2, KR1]. We denote by $\text{mod } \mathcal{C}$ the category of finitely presented $\mathcal{C}$ -modules. If $\mathcal{C}$ has pseudokernels, then $\text{mod } \mathcal{C}$ forms an abelian category [A1].

Proposition I.1.11. *Let $\mathcal{C}$ be an exact stably 2-CY category. Assume that $\mathcal{C}$ has pseudokernels and the global dimension of $\text{mod } \mathcal{C}$ is finite. Let $\Gamma = \text{End}(T)$ for a cluster tilting object T in $\mathcal{C}$.*

- (a) Γ has finite global dimension.
- (b) If $\mathcal{C}$ is Hom-finite, then the quiver of Γ has no loops. If moreover $\mathcal{C}$ is an extension closed subcategory of an abelian category closed under subobjects, then the quiver of Γ has no 2-cycles.

Proof. (a) Let $m = \text{gl.dim}(\text{mod } \mathcal{C})$. For any $X \in \text{mod } \Gamma$, take a projective presentation $(T, T_1) \rightarrow (T, T_0) \rightarrow X \rightarrow 0$. By our assumptions, there exists a complex $0 \rightarrow F_m \rightarrow \cdots \rightarrow F_2 \rightarrow T_1 \rightarrow T_0$ in $\mathcal{C}$ such that $0 \rightarrow (, F_m) \rightarrow \cdots \rightarrow (, F_2) \rightarrow (, T_1) \rightarrow (, T_0)$ is exact in $\text{mod } \mathcal{C}$. Since T is cluster tilting, we have an exact sequence $0 \rightarrow T_1 \rightarrow T_0 \rightarrow F_i \rightarrow 0$, with T_1 and T_0 in $\text{add } T$ by Proposition I.1.7. Hence we have $\text{pd}_\Gamma(T, F_i) \leq 1$ and consequently $\text{pd}_\Gamma X \leq m + 1$. It follows that Γ has finite global dimension.

(b) By (a), Γ is a finite dimensional algebra of finite global dimension. By [Le, Ig], the quiver of Γ has no loops.

We shall show the second assertion. Our proof is based on [GLS1, 6.4]. We start with showing that $\text{Ext}_\Gamma^2(S, S) = 0$ for any simple Γ -module S , assumed to be the top of the projective Γ -module (T, M) for an indecomposable summand M of T .

First, we assume that M is not projective in $\mathcal{C}$. Take exact exchange sequences $0 \rightarrow M^* \xrightarrow{f} B \xrightarrow{g} M \rightarrow 0$ and $0 \rightarrow M \xrightarrow{s} B' \xrightarrow{t} M^* \rightarrow 0$. Since Γ has no loops, we have a projective presentation $0 \rightarrow (T, M) \xrightarrow{s} (T, B') \xrightarrow{tf} (T, B) \xrightarrow{g} (T, M) \rightarrow S \rightarrow 0$ of the Γ -module S . Since M is not a summand of B' , we have $\text{Ext}_\Gamma^2(S, S) = 0$.

Next, we assume that M is projective in $\mathcal{C}$. Take a minimal projective presentation $(T, B) \xrightarrow{g} (T, M) \rightarrow S \rightarrow 0$ of the Γ -module S . By assumption, $\text{Im } g$ in the abelian category belongs to $\mathcal{C}$. Then $g: B \rightarrow \text{Im } g$ is a minimal right $\text{add } T$ -approximation. By Proposition I.1.7(a), we have that $B' = \text{Ker } g$ belongs to $\text{add } T$. Thus we have a projective resolution $0 \rightarrow (T, B') \rightarrow (T, B) \xrightarrow{g} (T, M) \rightarrow S \rightarrow 0$ of the Γ -module S . Since g is right minimal, B' does not have an injective summand. Thus we have $\text{Ext}_\Gamma^2(S, S) = 0$.

Since in both cases $\text{Ext}_\Gamma^2(S, S) = 0$, we can not have a 2-cycle by [GLS1, 3.11]. $\square$

I.2. Substructures.

For extension closed subcategories of triangulated or exact categories both having a weak cluster structure, we introduce the notion of substructure. Using heavily [IY], we give sufficient conditions for having a substructure, when starting with a Hom-finite triangulated 2-CY category or an exact stably 2-CY category, and using the cluster tilting subcategories, with the indecomposable projectives as coefficients.

Let $\mathcal{C}$ be an exact or triangulated k -category, and $\mathcal{B}$ a subcategory of $\mathcal{C}$ closed under extensions. Assume that both $\mathcal{C}$ and $\mathcal{B}$ have a weak cluster structure. We say that we have a *substructure* of $\mathcal{C}$ induced by an extended cluster T in $\mathcal{B}$ if we have the following:

There is a set A of indecomposable objects in $\mathcal{C}$ such that $\widetilde{T'} = T' \cup A$ is an extended cluster in $\mathcal{C}$ for any extended cluster T' in $\mathcal{B}$ which is obtained by a finite number of exchanges from T .

Note that for each sequence of cluster variables $M_1, \dots, M_t$, with M_{i+1} in $\mu_{M_i}(T)$, we have $\mu_{M_t}(\dots \mu_{M_1}(T)) \cup A = \tilde{\mu}_{M_t}(\dots \tilde{\mu}_{M_1}(\widetilde{T}))$, where μ denotes the exchange for $\mathcal{B}$ and $\tilde{\mu}$ the exchange for $\mathcal{C}$.

We shall investigate substructures arising from certain extension closed subcategories of triangulated 2-CY categories and of exact stably 2-CY categories. We start with the triangulated case, and here we first recall some results from [IY] specialized to the setting of 2-CY categories.

For a triangulated category $\mathcal{C}$ and full subcategories $\mathcal{B}$ and $\mathcal{B}'$, let $\mathcal{B}^\perp = \{X \in \mathcal{C} \mid \text{Hom}(\mathcal{B}, X) = 0\}$ and ${}^\perp\mathcal{B} = \{X \in \mathcal{C} \mid \text{Hom}(X, \mathcal{B}) = 0\}$. We denote by $\mathcal{B} * \mathcal{B}'$ the full subcategory of $\mathcal{C}$ consisting of all $X \in \mathcal{C}$ such that there exists a triangle $B \rightarrow X \rightarrow B' \rightarrow B[1]$ with $B \in \mathcal{B}$ and $B' \in \mathcal{B}'$.

We get the following sufficient conditions for constructing 2-CY categories, and hence categories with weak cluster structures.

Theorem I.2.1. *Let $\mathcal{C}$ be a Hom-finite triangulated 2-CY category and $\mathcal{B}$ a functorially finite extension closed subcategory of $\mathcal{C}$.*

- (a) $\mathcal{B}^\perp$ and ${}^\perp\mathcal{B}$ are functorially finite extension closed subcategories of $\mathcal{C}$. Moreover, $\mathcal{B} * \mathcal{B}^\perp = \mathcal{C} = {}^\perp\mathcal{B} * \mathcal{B}$ and ${}^\perp(\mathcal{B}^\perp) = \mathcal{B} = ({}^\perp\mathcal{B})^\perp$ hold.
- (b) Let $\mathcal{D} = \mathcal{B} \cap {}^\perp\mathcal{B}[1]$. Then $\mathcal{B}/\mathcal{D}$ is a Hom-finite triangulated 2-CY category, so that $\mathcal{B}$ is a stably 2-CY category. Moreover, $\mathcal{B} \subseteq (\mathcal{D} * \mathcal{B}[1]) \cap (\mathcal{B}[-1] * \mathcal{D})$ holds, and $\mathcal{D}$ is a functorially finite rigid subcategory of $\mathcal{C}$.
- (c) Let $\mathcal{D}$ be a functorially finite rigid subcategory of $\mathcal{C}$ and $\mathcal{B}' = {}^\perp\mathcal{D}[1]$. Then $\mathcal{B}'$ is a functorially finite extension closed subcategory of $\mathcal{C}$ and $\mathcal{B}'/\mathcal{D}$ is a triangulated 2-CY category. Moreover, there exists a one-one correspondence between cluster tilting (respectively, maximal rigid, rigid) subcategories of $\mathcal{C}$ containing $\mathcal{D}$ and cluster tilting (respectively, maximal rigid, rigid) subcategories of $\mathcal{B}'/\mathcal{D}$. It is given by $T \mapsto T/\mathcal{D}$.

Proof. (a) Since $\mathcal{B}^\perp = {}^\perp\mathcal{B}[2]$ holds by the 2-CY property, the assertion follows from [IY, 2.3].

(b) Clearly $\mathcal{B}/\mathcal{D}$ is Hom-finite, since $\mathcal{C}$ is. To show that $\mathcal{B}/\mathcal{D}$ is a triangulated 2-CY category, we only need to check $\mathcal{B} \subseteq (\mathcal{D} * \mathcal{B}[1]) \cap (\mathcal{B}[-1] * \mathcal{D})$ by [IY, 4.2]. Let Z be in $\mathcal{B}$. Since $\mathcal{B}$, and hence $\mathcal{B}[1]$, is functorially finite in $\mathcal{C}$, it follows from (a) that we have a triangle $X \rightarrow Y \rightarrow Z \rightarrow X[1]$ with Y in ${}^\perp\mathcal{B}[1]$ and $X[1]$ in $\mathcal{B}[1]$. Since $\mathcal{B}$ is extension closed, Y is in $\mathcal{B}$, and consequently Y is in $\mathcal{B} \cap {}^\perp\mathcal{B}[1] = \mathcal{D}$. It follows that Z is in $\mathcal{D} * \mathcal{B}[1]$, and similarly in $\mathcal{B}[-1] * \mathcal{D}$.

To see that $\mathcal{D}$ is functorially finite in $\mathcal{C}$, we only have to show that $\mathcal{D}$ is functorially finite in $\mathcal{B}$. For any $Z \in \mathcal{B}$, take the above triangle $X \rightarrow Y \xrightarrow{f} Z \rightarrow X[1]$ with Y in $\mathcal{D}$ and $X[1]$ in $\mathcal{B}[1]$. Since $(\mathcal{D}, X[1]) = 0$, we have that f is a right $\mathcal{D}$ -approximation. Thus $\mathcal{D}$ is contravariantly finite in $\mathcal{B}$, and similarly covariantly finite in $\mathcal{B}$.

(c) See [IY, 4.9]. □

The example of the cluster category $\mathcal{C}$ of the path algebra kQ where Q is of type A_3 from the previous section illustrates part of this theorem. For let $\mathcal{D} = \text{add } P_1$. Then $\mathcal{B}' = {}^\perp\mathcal{D}[1] = \text{mod } kQ$, and $\mathcal{B}'/\mathcal{D} = \mathcal{C}_{kQ'}$, where Q' is a quiver of type A_2 . The cluster tilting objects in $\mathcal{C}$ containing P_1 are $P_1 \oplus S_3 \oplus P_2$, $P_1 \oplus P_2 \oplus S_2$, $P_1 \oplus S_2 \oplus P_1/S_3$, $P_1 \oplus P_1/S_3 \oplus S_1$, $P_1 \oplus S_1 \oplus S_3$, which are in one-one correspondence with the cluster tilting objects in $\mathcal{B}'/\mathcal{D}$.

In order to get sufficient conditions for having a substructure we investigate cluster tilting subcategories in $\mathcal{B}$. For this the following lemma is useful.

Lemma I.2.2. *Let $\mathcal{C}$ be a Hom-finite triangulated 2-CY category. For any functorially finite and thick subcategory $\mathcal{C}_1$ of $\mathcal{C}$, there exists a functorially finite and thick subcategory $\mathcal{C}_2$ of $\mathcal{C}$ such that $\mathcal{C} = \mathcal{C}_1 \times \mathcal{C}_2$.*

Proof. Let $\mathcal{C}_2 = \mathcal{C}_1^\perp$. Then we have $\mathcal{C}_2 = \mathcal{C}_1^\perp = {}^\perp \mathcal{C}_1[2] = {}^\perp \mathcal{C}_1$ by Serre duality, using that $\mathcal{C}_1$ is triangulated. We only have to show that any object in $\mathcal{C}$ is a direct sum of objects in $\mathcal{C}_1$ and $\mathcal{C}_2$. For any $X \in \mathcal{C}$, there exists a triangle $A_1 \rightarrow X \rightarrow A_2 \xrightarrow{f} A_1[1]$ in $\mathcal{C}$ with A_1 in $\mathcal{C}_1$ and A_2 in $\mathcal{C}_2 = \mathcal{C}_1^\perp$ by Theorem I.2.1(a). Since $f = 0$, we have $X \simeq A_1 \oplus A_2$. Thus we have $\mathcal{C} = \mathcal{C}_1 \times \mathcal{C}_2$. $\square$

Using Lemma I.2.2, we get the following decomposition of triangulated categories.

Proposition I.2.3. *Let $\mathcal{C}$ be a Hom-finite triangulated 2-CY category and $\mathcal{B}$ a functorially finite extension closed subcategory of $\mathcal{C}$. Let $\mathcal{D} = \mathcal{B} \cap {}^\perp \mathcal{B}[1]$ and $\mathcal{B}' = {}^\perp \mathcal{D}[1]$.*

- (a) *There exists a functorially finite and extension closed subcategory $\mathcal{B}''$ of $\mathcal{C}$ such that $\mathcal{D} \subseteq \mathcal{B}'' \subseteq \mathcal{B}'$ and $\mathcal{B}' / \mathcal{D} = \mathcal{B} / \mathcal{D} \times \mathcal{B}'' / \mathcal{D}$ as a triangulated category.*
- (b) *There exists a one-one correspondence between pairs consisting of cluster tilting (respectively, maximal rigid, rigid) subcategories of $\mathcal{B}$ and of $\mathcal{B}''$, and cluster tilting (respectively, maximal rigid, rigid) subcategories of $\mathcal{B}'$. It is given by $(T, T'') \mapsto T \oplus T''$.*

Proof. (a) We know by Theorem I.2.1(b)(c) that $\mathcal{D}$ is functorially finite rigid, and that $\mathcal{B} / \mathcal{D}$ and $\mathcal{B}' / \mathcal{D}$ are both triangulated 2-CY categories. The inclusion functor $\mathcal{B} / \mathcal{D} \rightarrow \mathcal{B}' / \mathcal{D}$ is a triangle functor by the construction of their triangulated structures in [IY, 4.2]. In particular $\mathcal{B} / \mathcal{D}$ is a thick subcategory of $\mathcal{B}' / \mathcal{D}$, and hence we have a decomposition by Lemma I.2.2.

(b) This follows by Theorem I.2.1(c). $\square$

Then we get the following.

Corollary I.2.4. *Let $\mathcal{C}$ be a Hom-finite 2-CY algebraic triangulated category with a cluster tilting object, and $\mathcal{B}$ a functorially finite extension closed subcategory of $\mathcal{C}$. Then we have the following.*

- (a) *The stably 2-CY category $\mathcal{B}$ also has some cluster tilting object. Any maximal rigid object in $\mathcal{B}$ is a cluster tilting object in $\mathcal{B}$.*
- (b) *There is some rigid object A in $\mathcal{C}$ such that $T \oplus A$ is a cluster tilting object in $\mathcal{C}$ for any cluster tilting object T in $\mathcal{B}$.*
- (c) *Any cluster tilting object T in $\mathcal{B}$ determines a substructure for the weak cluster structures on $\mathcal{B}$ and $\mathcal{C}$ given by cluster tilting objects.*

Proof. (a) Let $\mathcal{D} = {}^\perp \mathcal{B}[1]$ and $\mathcal{B}' = {}^\perp \mathcal{D}[1]$. Since $\mathcal{C}$ is algebraic by Theorem I.1.8, we have a cluster tilting object T in $\mathcal{C}$ containing $\mathcal{D}$. By Proposition I.2.3, we have decompositions $\mathcal{B}' / \mathcal{D} = \mathcal{B} / \mathcal{D} \times \mathcal{B}'' / \mathcal{D}$ for some subcategory $\mathcal{B}''$ of $\mathcal{B}'$ and $T = T_1 \oplus T_2$ with a cluster tilting object T_1 (respectively, T_2) in $\mathcal{B}$ (respectively, $\mathcal{B}''$). Thus $\mathcal{B}$ has a cluster tilting object.

Now we show the second assertion. Let M be maximal rigid in $\mathcal{B}$. By Proposition I.2.3(b), we have that $M \oplus T_2$ is maximal rigid in $\mathcal{C}$. By Theorem I.1.8, it follows that $M \oplus T_2$ is cluster tilting in $\mathcal{C}$ and by Proposition I.2.3(b), we have that M is cluster tilting in $\mathcal{B}$.

(b) We only have to let $A = T_2$.

(c) This follows from (b). $\square$

It is curious to note that combining Proposition I.2.3 with Theorem I.2.1 we obtain a kind of classification of functorially finite extension closed subcategories of a triangulated 2-CY category in terms of functorially finite rigid subcategories, analogous to results from [AR1].

Theorem I.2.5. *Let $\mathcal{C}$ be a 2-CY triangulated category. Then the functorially finite extension closed subcategories $\mathcal{B}$ of $\mathcal{C}$ are all obtained as preimages under the functor $\pi: \mathcal{C} \rightarrow \mathcal{C} / \mathcal{D}$*

of the direct summands of ${}^\perp \mathcal{D}[1]/\mathcal{D}$ as a triangulated category, for functorially finite rigid subcategories $\mathcal{D}$ of $\mathcal{C}$.

Proof. Let $\mathcal{D}$ be functorially finite rigid in $\mathcal{C}$. Then $\mathcal{B}' = {}^\perp \mathcal{D}[1]$ is functorially finite extension closed in $\mathcal{C}$ by Theorem I.2.1(a). Then the preimage under $\pi: \mathcal{C} \rightarrow \mathcal{C}/\mathcal{D}$ of any direct summand of $\mathcal{B}'/\mathcal{D}$ as a triangulated category is functorially finite and extension closed in $\mathcal{C}$.

Conversely, let $\mathcal{B}$ be a functorially finite extension closed subcategory of $\mathcal{C}$ and $\mathcal{D} = \mathcal{B} \cap {}^\perp \mathcal{B}[1]$. By Proposition I.2.3, we have that $\mathcal{B}/\mathcal{D}$ is a direct summand of ${}^\perp \mathcal{D}[1]/\mathcal{D}$. $\square$

We now investigate substructures also for exact categories which are stably 2-CY. We have the following main result.

Theorem I.2.6. *Let $\mathcal{C}$ be an exact stably 2-CY category, and $\mathcal{B}$ a functorially finite extension closed subcategory of $\mathcal{C}$. Then $\mathcal{B}$ has enough projectives and injectives, and is a stably 2-CY category.*

Proof. We know that $\mathcal{B}$ is an exact category and $\mathcal{D} = \mathcal{B} \cap {}^\perp \mathcal{B}[1]$ is the subcategory of projective injective objects. Since $\mathcal{B} \subseteq \mathcal{B}[-1] * \mathcal{D}$ holds by Theorem I.2.1(b), then for any $X \in \mathcal{B}$, there exists a triangle $X \rightarrow Y \rightarrow Z \rightarrow X[1]$ with $Y \in \mathcal{D}$ and $Z \in \mathcal{B}$. This is induced from an exact sequence $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$ in $\mathcal{C}$. Thus $\mathcal{B}$ has enough injectives. Dually, $\mathcal{B}$ has enough projectives, which coincide with the injectives. Hence $\mathcal{B}$ is a Frobenius category, and consequently, $\mathcal{B}$ is stably 2-CY. $\square$

Alternatively we give a direct approach, where the essential information is given by the following lemma and its dual.

Lemma I.2.7. *Let $\mathcal{C}$ be an exact category with enough injectives, and $\mathcal{B}$ a contravariantly finite extension closed subcategory of $\mathcal{C}$. Then $\mathcal{B}$ is an exact category with enough injectives.*

Proof. It is clear that $\mathcal{B}$ is also an exact category. Let X be in $\mathcal{C}$ and take an exact sequence $0 \rightarrow X \rightarrow I \rightarrow X' \rightarrow 0$ with I injective in $\mathcal{C}$. Then we have an exact sequence of functors $(\cdot, X') \rightarrow \text{Ext}^1(\cdot, X) \rightarrow 0$. Since $\mathcal{B}$ is Krull-Schmidt and contravariantly finite in $\mathcal{C}$, we can take a projective cover $\phi: (\cdot, Y) \rightarrow \text{Ext}^1(\cdot, X)|_{\mathcal{B}} \rightarrow 0$ of $\mathcal{B}$ -modules. This is induced by an exact sequence $0 \rightarrow X \rightarrow Z \rightarrow Y \rightarrow 0$ with terms in $\mathcal{B}$.

We will show that Z is injective. Take any exact sequence $0 \rightarrow Z \rightarrow Z' \rightarrow Z'' \rightarrow 0$ with terms in $\mathcal{B}$. We will show that this splits. Consider the following exact commutative diagram:

$$\begin{array}{ccccccccc}
 & & & & 0 & & 0 & & \\
 & & & & \downarrow & & \downarrow & & \\
 0 & \rightarrow & X & \rightarrow & Z & \rightarrow & Y & \rightarrow & 0 \\
 & & \parallel & & \downarrow & & \downarrow^a & & \\
 0 & \rightarrow & X & \rightarrow & Z' & \rightarrow & Y' & \rightarrow & 0 \\
 & & & & \downarrow & & \downarrow & & \\
 & & & & Z'' & = & Z'' & & \\
 & & & & \downarrow & & \downarrow & & \\
 & & & & 0 & & 0 & &
 \end{array}
 \tag{2}$$

Then $Y' \in \mathcal{B}$, and we have the commutative diagram

$$\begin{array}{ccccccccc}
 0 & \rightarrow & (\cdot, X) & \rightarrow & (\cdot, Z) & \rightarrow & (\cdot, Y) & \xrightarrow{\phi} & \text{Ext}^1(\cdot, X)|_{\mathcal{B}} \rightarrow 0 \\
 & & \parallel & & \downarrow & & \downarrow^a & & \parallel \\
 0 & \rightarrow & (\cdot, X) & \rightarrow & (\cdot, Z') & \rightarrow & (\cdot, Y') & \rightarrow & \text{Ext}^1(\cdot, X)|_{\mathcal{B}}
 \end{array}
 \tag{3}$$

of exact sequences of $\mathcal{B}$ -modules. Since ϕ is a projective cover, we have that $(\cdot a)$ is a split monomorphism. Thus a is a split monomorphism. We see that the sequence $0 \rightarrow \text{Ext}^1(Z'', Z) \rightarrow \text{Ext}^1(Z'', Y)$ is exact by evaluating the upper sequence in (3) at Z'' . Since

the right vertical sequence in (2) splits, it follows that the middle vertical sequence in (2) splits. Hence Z is injective, and consequently $\mathcal{B}$ has enough injectives. $\square$

It follows from $\mathcal{C}$ being 2-CY that the projectives and injectives in $\mathcal{B}$ coincide, and hence $\mathcal{B}$ is Frobenius by Lemma I.2.7 and its dual. It follows as before that $\mathcal{B}$ is stably 2-CY, and the alternative proof of Theorem I.2.6 is completed.

We have the following interesting special case, as a consequence of Theorem I.2.6 and Corollary I.2.4. For X in $\mathcal{C}$ we denote by $\text{Sub } X$ the subcategory of $\mathcal{C}$ whose objects are subobjects of finite direct sums of copies of X .

Corollary I.2.8. *Let $\mathcal{C}$ be a Hom-finite abelian stably 2-CY category, and let X be an object in $\mathcal{C}$ with $\text{Ext}^1(X, X) = 0$, and $\text{id } X \leq 1$.*

- (a) *Then $\text{Sub } X$ is a functorially finite extension closed subcategory of $\mathcal{C}$ and is exact stably 2-CY.*
- (b) *If $\mathcal{C}$ has a cluster tilting object, then so does $\text{Sub } X$, and any cluster tilting object in $\text{Sub } X$ determines a substructure of the cluster structure for $\mathcal{C}$.*
- (c) *If $\mathcal{C}$ is abelian, then $\text{Sub } X$ has no loops or 2-cycles.*

Proof. (a) We include the proof for the convenience of the reader. We first want to show that $\text{Sub } X$ is extension closed. Let $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ be an exact sequence with A and C in $\text{Sub } X$, and consider the diagram

$$\begin{array}{ccccccc}
 & & 0 & & & 0 & \\
 & & \downarrow & & & \downarrow & \\
 0 & \longrightarrow & A & \xrightarrow{i} & B & \xrightarrow{j} & C \longrightarrow 0 \\
 & & \downarrow f & & & & \downarrow q \\
 0 & \longrightarrow & X_0 & \longrightarrow & X_0 \oplus X_1 & \longrightarrow & X_1 \longrightarrow 0
 \end{array}$$

with X_0, X_1 in $\text{add } X$. Since $\text{id } X_0 \leq 1$, we have the exact sequence $\text{Ext}^1(X_1, X_0) \rightarrow \text{Ext}^1(C, X_0) \rightarrow 0$, which shows that $\text{Ext}^1(C, X_0) = 0$. Then the exact sequence $(B, X_0) \rightarrow (A, X_0) \rightarrow \text{Ext}^1(C, X_0)$ shows that there is some $t: B \rightarrow X_0$ such that $it = f$. This shows that B is in $\text{Sub } X$, which is then closed under extensions. It is also functorially finite [AS], and clearly Krull-Schmidt. So $\text{Sub } X$ has enough projectives and injectives, with the projectives coinciding with the injectives. Hence $\text{Sub } X$ is Frobenius, and as we have seen before, it follows that the stable category $\underline{\text{Sub } X}$ is 2-CY.

(b) This follows directly using Corollary I.2.4.

(c) This follows from Proposition I.1.11. $\square$

In order to see when we have cluster structures we next want to give sufficient conditions for algebraic triangulated (or stably) 2-CY categories not to have loops or 2-cycles.

Proposition I.2.9. *Let $\mathcal{C}$ be a Hom-finite algebraic triangulated (or exact stably) 2-CY category with a cluster tilting object, and $\mathcal{B}$ a functorially finite extension closed subcategory.*

- (a) *If $\mathcal{C}$ has no 2-cycles, then also $\mathcal{B}$ has no 2-cycles.*
- (b) *If $\mathcal{C}$ has no loops, then $\mathcal{B}$ has no loops.*

Proof. We give a proof for the algebraic triangulated 2-CY case. A similar argument works for the stably 2-CY case.

(a) Let $\mathcal{D} = \mathcal{B} \cap {}^\perp \mathcal{B}[1]$ and $\mathcal{B}' = {}^\perp \mathcal{D}[1]$. Since cluster tilting objects in $\mathcal{B}'$ are exactly cluster tilting objects in $\mathcal{C}$ which contain $\mathcal{D}$, our assumption implies that $\mathcal{B}'$ has no 2-cycles.

We shall show that $\mathcal{B}$ has no 2-cycles. Let T be a cluster tilting object in $\mathcal{B}$. By Corollary I.2.4(b), there exists $T' \in \mathcal{B}'$ such that $T \oplus T'$ is a cluster tilting object in $\mathcal{C}$. We already observed that $T \oplus T'$ has no 2-cycles. If T has a 2-cycle, then at least one arrow in the

2-cycle represents a morphism $f: X \rightarrow Y$ which factors through an object in T' . We write f as a composition of $f_1: X \rightarrow Z$ and $f_2: Z \rightarrow Y$ with $Z \in T'$. Since $\mathcal{B}/\mathcal{D}$ is a direct summand of $\mathcal{B}'/\mathcal{D}$ by Proposition I.2.3, any morphism between T and T' factors through $\mathcal{D}$. Thus we can write f_1 (respectively f_2) as a composition of $g_1: X \rightarrow W_1$ and $h_1: W_1 \rightarrow Z$ (respectively, $g_2: Z \rightarrow W_2$ and $h_2: W_2 \rightarrow Y$) with $W_1 \in \mathcal{D}$ (respectively, $W_2 \in \mathcal{D}$). We have $f = f_1 f_2 = g_1(h_1 g_2)h_2$, where $h_1 g_2$ is in $\text{rad } \mathcal{B}$ and at least one of h_2 and g_1 is in $\text{rad } \mathcal{B}$, since at least one of X and Y is not in $\mathcal{D}$. So f can not be irreducible in $\text{add } T$, a contradiction.

(b) This follows in a similar way. $\square$

Note that the quiver Q_T may have 2-cycles between coefficients. For example, let $\mathcal{C} = \text{mod } \Lambda$ for the preprojective algebra of a Dynkin quiver and let $\mathcal{B}$ be the subcategory $\text{add } \Lambda$. Then there are no 2-cycles for $\mathcal{C}$, but there are 2-cycles for $\mathcal{B}$, since Λ is the only cluster tilting object in $\mathcal{B}$.

I.3. Preprojective algebras of Dynkin type.

In this section we specialize our general results from Section I.2 to the case of the finitely generated modules over a preprojective algebra of Dynkin type. We also illustrate with three concrete examples. The same examples will be used in the Chapter III to illustrate how to use this theory to construct subcluster algebras of cluster algebras.

The category $\mathcal{C} = \text{mod } \Lambda$ for Λ preprojective of Dynkin type is a Hom-finite Frobenius category. By [GLS1], see also Section II.2, a rigid Λ -module is cluster tilting if and only if the number of non-isomorphic indecomposable summands is the number of positive roots, so $\frac{n(n+1)}{2}$ for A_n , $n(n-1)$ for D_n , 36 for E_6 , 63 for E_7 and 120 for E_8 .

Let $\mathcal{B}$ be an extension closed functorially finite subcategory of $\mathcal{C}$. We know that $\mathcal{B}$ is stably 2-CY by Theorem I.2.6. It is known that $\mathcal{C}$ and $\underline{\mathcal{C}}$ have no loops or 2-cycles for the cluster tilting objects [GLS1], and this also follows from Proposition I.1.11. Then it follows from Proposition I.2.9 that there are also no loops or 2-cycles for $\mathcal{B}$ and the subcategory $\underline{\mathcal{B}}$ of $\underline{\mathcal{C}}$. Note that $\underline{\mathcal{B}}$ is not the stable category of $\mathcal{B}$ since $\mathcal{B}$ may have more projectives than $\mathcal{C}$.

We then have the following.

Theorem I.3.1. *Let $\mathcal{B}$ be an extension closed functorially finite subcategory of the category $\mathcal{C} = \text{mod } \Lambda$ for the preprojective algebra Λ of a Dynkin quiver. Then we have the following.*

- (a) *The exact stably 2-CY category $\mathcal{B}$ has a cluster tilting object, and any maximal rigid object in $\mathcal{B}$ is a cluster tilting object, which can be extended to a cluster tilting object for $\mathcal{C}$, and which gives rise to a substructure.*
- (b) *The category $\underline{\mathcal{B}}$ is a stably 2-CY Frobenius category with no loops or 2-cycles for the cluster tilting objects, and hence has a cluster structure.*

Proof. (a) This follows from Theorem I.1.8 and Corollary I.2.8.

(b) This follows from the above comments and Theorem I.1.6. $\square$

We now give some concrete examples of weak cluster structures and substructures. In Chapter III these examples will be revisited, and used to model cluster algebras and subcluster algebras.

We denote by P_i the indecomposable projective module associated to vertex i , by J the radical of a ring, and by S_i the simple top of P_i . Usually, we represent a module M by its radical filtration, the numbers in the first row represent the indices of the simples in M/JM , and the numbers in the i 'th row represent the indices of the simples in $J^{i-1}M/J^iM$. e.g.

$$\begin{array}{ccc} & 2 & \\ 1 & & 3 \\ & 2 & \end{array}$$

represents the indecomposable projective module P_2 for the preprojective algebra of type A_3 , which has quiver

$$1 \rightleftarrows 2 \rightleftarrows 3$$


Example 1. Let Λ be the preprojective algebra of a Dynkin quiver A_4 . This algebra has quiver

$$1 \rightleftarrows 2 \rightleftarrows 3 \rightleftarrows 4$$

We consider the modules P_3 and $M = JP_3$. These modules are represented by their radical filtrations:

$$\begin{smallmatrix} & 3 \\ 1 & 2 & 3 & 4 \\ & 2 \end{smallmatrix} \quad \text{and} \quad \begin{smallmatrix} & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \\ & 2 \end{smallmatrix}$$

We let $\mathcal{C}' = \text{Sub } P_3$ and $\mathcal{B} = \{X \in \mathcal{C}' \mid \text{Ext}^1(M, X) = 0\}$. The AR-quiver of $\mathcal{C}'$ is given below, where we name the indecomposables in $\mathcal{C}'$ to ease notation. The indexing will be explained in Section III.2.



From the AR-quiver we see that the indecomposable projectives in $\mathcal{C}'$ are M_{45}, M_{34}, M_{23} and M_{15} . The indecomposables of the subcategory $\mathcal{B}$ are obtained from $\mathcal{C}'$ by deleting the indecomposable objects $P_3 = M_{45}, M_x$ and M_y . The category $\mathcal{B}$ is extension closed by definition, and the indecomposable projectives in $\mathcal{B}$ are $M_{35}, M_{34}, M_{23}, M_{15}$.

Let $T = M_{34} \oplus M_{23} \oplus M_{13} \oplus M_{15} \oplus M_{35}$, then clearly $\text{Ext}^1(T, T) = 0$, and the unique indecomposable in $\mathcal{B}$ which is not a summand in T is M_{25} , which has a non-zero extension with T . Hence T is a cluster tilting object in $\mathcal{B}$, and $\mathcal{B}$ has a cluster structure, with coefficients $M_{35}, M_{34}, M_{23}, M_{15}$.

Now, since M_{45} is projective in $\mathcal{C}'$, and M_x as well as M_y has non-zero extensions with T , it is clear that $T' = T \oplus M_{45}$ is a cluster tilting object in $\mathcal{C}'$, and hence $\mathcal{C}'$ has a cluster structure, such that we have a substructure for $\mathcal{B}$ induced by T .

We claim that the cluster tilting object T' in $\mathcal{C}'$ can be extended to a cluster tilting object $\tilde{T} = T' \oplus P_1 \oplus P_2 \oplus P_4 \oplus Z$ of $\text{mod } \Lambda$, where Z is the Λ -module with radical filtration $\begin{smallmatrix} 1 \\ 2 \\ 3 \end{smallmatrix}$.

To see that $\text{Ext}^1(\tilde{T}, \tilde{T}) = 0$, it is sufficient to show that $\text{Ext}^1(Z, X \oplus Z) = 0$ for all X in $\mathcal{C}'$. There is an exact sequence $0 \rightarrow S_4 \rightarrow P_1 \rightarrow Z \rightarrow 0$, and hence for every X in $\mathcal{C}'$, there is an exact sequence


$$\text{Hom}(S_4, X \oplus Z) \rightarrow \text{Ext}^1(Z, X \oplus Z) \rightarrow \text{Ext}^1(P_1, X \oplus Z).$$

Note that $\text{Hom}(S_4, P_3) = 0$ and hence $\text{Hom}(S_4, X) = 0$ for all X in $\mathcal{C}'$, and that $\text{Hom}(S_4, Z) = 0$. It follows that $\text{Ext}^1(Z, X \oplus Z) = 0$. Thus $\tilde{T}$ is a cluster tilting object since it has the correct number $10 = \frac{4 \cdot 5}{2}$ of indecomposable direct summands.

Example 2. For our next example, let Λ be the preprojective algebra of type A_3 . It has the quiver

$$1 \rightleftarrows 2 \rightleftarrows 3$$


The AR-quiver of $\mathcal{C} = \text{mod } \Lambda$ is given by the following. We name the indecomposables in $\mathcal{C}$ according to the following table. The indexing will be explained in Section III.2.



The indecomposable projectives in $\mathcal{C}$ are M_{34}, M_z, M_{234} .


Let $\mathcal{B}$ be the full subcategory of $\mathcal{C}$ generated by $P_2 \oplus P_3$. Then $\mathcal{B} = \text{add}(M_{34} \oplus M_{124} \oplus M_{24} \oplus M_{23} \oplus M_{134} \oplus M_{234} \oplus M_{13} \oplus M_{14})$. In addition to M_{34}, M_{234} , also M_{134} becomes projective in $\mathcal{B}$. It is straightforward to see that $M_{23} \oplus M_{13}$ is extension-free, so $T = M_{34} \oplus M_{234} \oplus M_{134} \oplus M_{23} \oplus M_{13}$ has $\text{Ext}^1(T, T) = 0$. Let $\tilde{T} = T \oplus M_z$. Then also $\text{Ext}^1(\tilde{T}, \tilde{T}) = 0$. Since $\tilde{T}$ has the correct number of indecomposable direct summands $6 = \frac{3+4}{2}$, it is a cluster tilting object. Hence T is a cluster tilting object in $\mathcal{B}$.

Example 3. In this example we let Λ be the preprojective algebra of a Dynkin quiver D_4 .



We consider the subcategory $\mathcal{B} = \text{Sub } P_2$. Using Corollary I.2.8 we have that $\mathcal{B}$ is extension closed. We know by Theorem I.3.1 that $\mathcal{B}$ has a cluster tilting object that can be extended to a cluster tilting object for $\mathcal{C} = \text{mod } \Lambda$.

The following gives P_2 as a representation of the quiver with relations



The modules in $\mathcal{B}$ do not necessarily have a simple socle, and in fact the subcategory is not of finite type. As noted earlier, it is functorially finite. However, the indecomposable direct summands in the cluster tilting object we will construct all have simple socle.

The indecomposable submodules of P_2 we will need to construct a cluster tilting object have the following radical filtrations. The indexing will be explained in Chapter III.

M_{16}	$\begin{smallmatrix} & 3 & 4 \\ & 2 & \\ & & \end{smallmatrix}$	M_{24}	$\begin{smallmatrix} & 1 & 3 \\ & 2 & \\ & & \end{smallmatrix}$	M_{25}	$\begin{smallmatrix} & 1 & 4 \\ & 2 & \\ & & \end{smallmatrix}$	M_{26}	$\begin{smallmatrix} & 1 & 3 & 4 \\ & 2 & & \\ & & & \end{smallmatrix}$
M_{68}	$\begin{smallmatrix} & 1 & 3 & 4 \\ & 2 & 2 & \\ & 1 & 3 & 4 \\ & & 2 & \end{smallmatrix}$	M_{18}	$\begin{smallmatrix} & 1 & \\ & 2 & 4 \\ & 3 & 2 \\ & & 2 \end{smallmatrix}$	M_-	$\begin{smallmatrix} & 4 \\ & 2 & 3 \\ & 1 & 2 \\ & & 2 \end{smallmatrix}$	M_+	$\begin{smallmatrix} & 3 & 4 \\ & 2 & \\ & 1 & 2 & 4 \\ & & 2 & \end{smallmatrix}$

The indecomposable projectives in $\mathcal{B}$ are P_2, M_{18}, M_+ and M_- . This follows from the following.

Lemma I.3.2. *Let Λ be a finite dimensional algebra with X in $\text{mod } \Lambda$ such that $\text{Sub } X$ is extension closed in $\text{mod } \Lambda$. Then the indecomposable projective objects in $\text{Sub } X$ are of the form $P/\alpha(P)$ where P is an indecomposable projective Λ -module and $\alpha(P)$ is the smallest submodule of P such that $P/\alpha(P)$ is in $\text{Sub } X$.*

Proof. For convenience of the reader, we include a proof. Let P be indecomposable projective in $\text{mod } \Lambda$. It is clear that there is a smallest submodule $\alpha(P)$ of P such that $P/\alpha(P)$ is in $\text{Sub } X$. For, if A and B are submodules of P with P/A and P/B in $\text{Sub } X$, then clearly $P/A \cap B \subseteq P/A \oplus P/B$ is in $\text{Sub } X$. It is clear that the natural map $f: P \rightarrow P/\alpha(P)$ is a minimal left $\text{Sub } X$ -approximation and that every module in $\text{Sub } X$ is a factor of a direct sum of Λ -modules of the form $P/\alpha(P)$ for P indecomposable projective. To see that each $P/\alpha(P)$ is projective in $\text{Sub } X$, consider the exact sequence $0 \rightarrow A \xrightarrow{s} B \xrightarrow{t} P/\alpha(P) \rightarrow 0$ in $\text{Sub } X$. Then there is some $u: P \rightarrow B$ such that $ut = f$. Since B is in $\text{Sub } X$, then $u(\alpha(P)) = 0$, so the sequence splits. Clearly there are no other indecomposable projectives in $\text{Sub } X$ since all modules in $\text{Sub } X$ are factors of direct sums of those of the form $P/\alpha(P)$. $\square$

In addition we need the following, where we leave the details to the reader.

Lemma I.3.3. *Let $M = M_{16} \oplus M_{24} \oplus M_{25} \oplus M_{26} \oplus M_{68}$. Then we have $\text{Ext}^1(M, M) = 0$.*

Hence, the module $T = M \oplus P_2 \oplus M_{18} \oplus M_+ \oplus M_-$ in $\mathcal{B}$ is rigid. If we add the other projectives we obtain the module $\tilde{T} = T \oplus P_1 \oplus P_3 \oplus P_4$, which also satisfies $\text{Ext}^1(\tilde{T}, \tilde{T}) = 0$.

Since $\tilde{T}$ has the correct number $12 = 4 \cdot 3$ of indecomposable summands, it is a cluster tilting object in $\mathcal{C} = \text{mod } \Lambda$. It is also clear from this that T is a cluster tilting object in $\mathcal{B} = \text{Sub } P_2$, since we added only projectives/injectives to T to obtain $\tilde{T}$. Note that $\mathcal{B}$ has a substructure of the cluster structure of $\mathcal{C}$.

II. PREPROJECTIVE ALGEBRAS FOR NON-DYCKIN QUIVERS

In this chapter we deal with completions of preprojective algebras of a finite connected quiver Q with no oriented cycles, and mainly those which are not Dynkin. In this case the modules of finite length coincide with the nilpotent modules over the preprojective algebra. These algebras Λ are known to be derived 2-CY (see [B, CB, BBK, GLS4]). Tilting Λ -modules of projective dimension at most one were investigated in [IR] when the quiver Q is a (generalized) extended Dynkin quiver. It was shown that such tilting modules are exactly the ideals in Λ which are finite products of two-sided ideals $I_i = \Lambda(1 - e_i)\Lambda$, where $e_1, \dots, e_n$ correspond to the vertices of the quiver, and that they are in one-one correspondence with the elements of the corresponding Weyl group, where $w = s_{i_1} \cdots s_{i_k}$ corresponds to $I_w = I_{i_1} \cdots I_{i_k}$. Here we generalize some of the results from [IR] beyond the noetherian case. In particular, we show that any finite product of ideals of the form I_i is a tilting module, and show that there is a bijection between cofinite tilting ideals and elements of the associated Coxeter group W .

For any descending chain of tilting ideals of the form $\Lambda \supseteq I_{i_1} \supseteq I_{i_1} I_{i_2} \supseteq I_{i_1} I_{i_2} \cdots I_{i_k} \supseteq \cdots$ we show that for $\Lambda_m = \Lambda / I_{i_1} \cdots I_{i_m}$, the categories $\text{Sub } \Lambda_m$ and $\underline{\text{Sub}} \Lambda_m$ are respectively stably 2-CY and 2-CY with nice cluster tilting objects. In this way we get, for any $w \in W$, a stably 2-CY category $\mathcal{C}_w = \text{Sub}(\Lambda / I_w)$, and for any reduced expression $w = s_{i_1} \cdots s_{i_k}$, a cluster tilting object $\bigoplus_{j=1}^k \Lambda / I_{s_{i_1} \cdots s_{i_j}}$ in $\mathcal{C}_w$. We also construct cluster tilting subcategories of the derived 2-CY category f.l. Λ . This way we get many examples of weak cluster structures without loops or 2-cycles which are then cluster structures by Theorem I.1.6. We also get many examples of substructures. In particular, any cluster category and the stable category $\underline{\text{mod}} \Lambda$ of a preprojective algebra of Dynkin type occur amongst this class. We give a description of the quivers of the cluster tilting objects/subcategories in terms of the associated reduced expressions. For example, the quiver of the preprojective component of the hereditary algebra with additional arrows from X to τX occur this way. In Section III.3 results in this chapter are used to show that coordinate rings of some unipotent cells of $\text{SL}_2(\mathbb{C}[t, t^{-1}])$ have a cluster algebra structure.

We refer to [I3] for corresponding results for d -CY algebras.

II.1. Tilting modules over 2-CY algebras.

Let Q be a finite connected quiver without oriented cycles which is not Dynkin, k an algebraically closed field and Λ the completion of the associated preprojective algebra. In [IR] the tilting Λ -modules of projective dimension at most one were investigated in the noetherian case, that is, when Q is extended Dynkin [BGL] (and also the generalized ones having loops). In this section we generalize some of these results to the non-noetherian case, concentrating on the aspects that will be needed for our construction of new 2-CY categories with cluster tilting objects/subcategories in the next sections. Note that since Λ is complete, the Krull-Schmidt theorem holds for finitely generated projective Λ -modules.

We say that a finitely presented Λ -module T is a *tilting module* if (i) there exists an exact sequence $0 \rightarrow P_n \rightarrow \cdots \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$ with finitely generated projective Λ -modules P_i , (ii) $\text{Ext}_\Lambda^i(T, T) = 0$ for any $i > 0$, (iii) there exists an exact sequence $0 \rightarrow \Lambda \rightarrow T_0 \rightarrow \cdots \rightarrow T_n \rightarrow 0$ with T_i in $\text{add } T$.

We say that $T \in \mathbf{D}(\text{Mod } \Lambda)$ a *tilting complex* [Ri] if (i') T is quasi-isomorphic to an object in the category $\mathbf{K}^b(\text{pr } \Lambda)$ of bounded complexes of finitely generated projective Λ -modules $\text{pr } \Lambda$, (ii') $\text{Hom}_{\mathbf{D}(\text{Mod } \Lambda)}(T, T[i]) = 0$ for any $i \neq 0$, (iii') T generates $\mathbf{K}^b(\text{pr } \Lambda)$.

A tilting module is nothing but a module which is a tilting complex since the condition (iii) can be replaced by (iii'). A *partial tilting complex* is a direct summand of a tilting complex. A *partial tilting module* is a module which is a partial tilting complex.

Let $1, \dots, n$ denote the vertices in Q , and let $e_1, \dots, e_n$ be the corresponding idempotents. For each i we denote by I_i the ideal $\Lambda(1 - e_i)\Lambda$. Then $S_i = \Lambda/I_i$ is a simple Λ -module and Λ^{op} -module since by assumption there are no loops in the quiver. We shall show that each I_i , and any finite product of such ideals, is a tilting ideal in Λ , and give some information about how the different products are related. But first we give several preliminary results, where new proofs are needed compared to [IR] since we do not assume Λ to be noetherian.

Lemma II.1.1. *Let T be a partial tilting Λ -module of projective dimension at most 1 and S a simple Λ^{op} -module. Then at least one of the statements $S \otimes_{\Lambda} T = 0$ and $\text{Tor}_1^{\Lambda}(S, T) = 0$ holds.*

Proof. We only have to show that there is a projective resolution $0 \rightarrow P_1 \rightarrow P_0 \rightarrow T \rightarrow 0$ such that P_0 and P_1 do not have a common summand. This is shown as in [HU, 1.2]. $\square$

Recall that for rings Λ and Γ , we call an object T in $\mathbf{D}(\text{Mod } \Lambda \otimes_{\mathbb{Z}} \Gamma^{\text{op}})$ a *two-sided tilting complex* if T is a tilting complex in $\mathbf{D}(\text{Mod } \Lambda)$ and $\text{End}_{\mathbf{D}(\text{Mod } \Lambda)}(T) \simeq \Gamma$ naturally.

The following result is useful (see [Ri][Ye, 1.7]).

Lemma II.1.2. *Let $T \in \mathbf{D}(\text{Mod } \Lambda \otimes_{\mathbb{Z}} \Gamma^{\text{op}})$ be a two-sided tilting complex.*

- (a) *For any tilting complex (respectively, partial tilting complex) U of Γ , we have a tilting complex (respectively, partial tilting complex) $T \overset{\mathbf{L}}{\otimes}_{\Gamma} U$ of Λ such that $\text{End}_{\mathbf{D}(\text{Mod } \Lambda)}(T \overset{\mathbf{L}}{\otimes}_{\Gamma} U) \simeq \text{End}_{\mathbf{D}(\text{Mod } \Gamma)}(U)$.*
- (b) *$\mathbf{R}\text{Hom}_{\Lambda}(T, \Lambda)$ and $\mathbf{R}\text{Hom}_{\Gamma^{\text{op}}}(T, \Gamma)$ are two-sided tilting complexes and isomorphic in $\mathbf{D}(\text{Mod } \Gamma \otimes_{\mathbb{Z}} \Lambda^{\text{op}})$.*

We collect some basic information on preprojective algebras.

Proposition II.1.3. *Let Λ be the completion of the preprojective algebra of a finite connected non-Dynkin diagram without loops.*

- (a) *Let Γ be the completion of $\Lambda \otimes_{kQ_0} \Lambda^{\text{op}}$ with respect to the ideal $J \otimes_{kQ_0} \Lambda^{\text{op}} + \Lambda \otimes_{kQ_0} J^{\text{op}}$ where J is the radical of Λ . Then there exists a commutative diagram*

$$\begin{array}{ccccccccc} 0 & \rightarrow & P_2 & \xrightarrow{f_2} & P_1 & \xrightarrow{f_1} & P_0 & \rightarrow & \Lambda & \rightarrow & 0 \\ & & \downarrow \wr & & \downarrow \wr & & \downarrow \wr & & \downarrow \wr & & \\ 0 & \rightarrow & \text{Hom}_{\Gamma}(P_0, \Gamma) & \xrightarrow{f_1^*} & \text{Hom}_{\Gamma}(P_1, \Gamma) & \xrightarrow{f_2^*} & \text{Hom}_{\Gamma}(P_2, \Gamma) & \rightarrow & \Lambda & \rightarrow & 0 \end{array}$$

of exact sequences of Γ -modules such that each P_i is a finitely generated projective Γ -module and $P_0 \simeq P_2 \simeq \Gamma$.

- (b) *There exists a functorial isomorphism $\text{Hom}_{\mathbf{D}(\text{Mod } \Lambda)}(X, Y[1]) \simeq D \text{Hom}_{\mathbf{D}(\text{Mod } \Lambda)}(Y, X[1])$ for any $X \in \mathbf{D}^b(\text{f.l. } \Lambda)$ and $Y \in \mathbf{K}^b(\text{pr } \Lambda)$.*
- (c) *f.l. Λ is derived 2-CY and $\text{gl. dim } \Lambda = 2$. In particular, any left ideal I of Λ satisfies $\text{pd}_{\Lambda} I \leq 1$.*
- (d) *$\text{Ext}_{\Lambda}^i(X, \Lambda) = 0$ for $i \neq 2$ and $\text{Ext}_{\Lambda}^2(X, \Lambda) \simeq DX$ for any $X \in \text{f.l. } \Lambda$.*

Proof. (a) See [GLS1, Section 8] and [BBK, Section 4.1].

(b) This follow from (a) and [B, 4.2].

(c)(d) Immediate from (a) and (b). $\square$

We are now ready to show that each I_i , and a finite product of such ideals, is a tilting module.

Proposition II.1.4. I_i is a tilting Λ -module of projective dimension at most one and $\text{End}_\Lambda(I_i) = \Lambda$.

Proof. We have $\text{Ext}_\Lambda^n(S_i, \Lambda) \simeq D\text{Ext}_\Lambda^{2-n}(\Lambda, S_i) = 0$ for $n = 0, 1$ by Proposition II.1.3. Applying $\text{Hom}_\Lambda(-, \Lambda)$ to the exact sequence $0 \rightarrow I_i \rightarrow \Lambda \rightarrow S_i \rightarrow 0$, we get $\text{Hom}_\Lambda(I_i, \Lambda) = \Lambda$. Applying $\text{Hom}_\Lambda(I_i, -)$, we get an exact sequence $0 \rightarrow \text{End}_\Lambda(I_i) \rightarrow \text{Hom}_\Lambda(I_i, \Lambda) \rightarrow \text{Hom}_\Lambda(I_i, S_i)$. Since $\text{Hom}_\Lambda(I_i, S_i) = 0$, we have $\text{End}_\Lambda(I_i) = \text{Hom}_\Lambda(I_i, \Lambda) = \Lambda$.

Applying $\otimes_\Lambda S_i$ to the exact sequence in Proposition II.1.3(a) we have a projective resolution

$$(4) \quad 0 \rightarrow \Lambda e_i \xrightarrow{g} P \xrightarrow{f} \Lambda e_i \rightarrow S_i \rightarrow 0$$

with $\text{Im } f = I_i e_i$ and $P \in \text{add } \Lambda(1 - e_i)$. In particular $I_i = \text{Im } f \oplus \Lambda(1 - e_i)$ is a finitely presented Λ -module with $\text{pd } I_i \leq 1$.

We have $\text{Ext}_\Lambda^1(I_i, I_i) \simeq \text{Ext}_\Lambda^2(S_i, I_i) \simeq D\text{Hom}_\Lambda(I_i, S_i) = 0$. Using (4), we have an exact sequence

$$0 \rightarrow \Lambda \rightarrow P \oplus \Lambda(1 - e_i) \rightarrow I_i e_i \rightarrow 0$$

such that the middle and the right terms belong to $\text{add } I_i$. Thus I_i is a tilting Λ -module. $\square$

Proposition II.1.5. Let T be a tilting Λ -module of projective dimension at most one.

- (a) If $\text{Tor}_1^\Lambda(S_i, T) = 0$, then $I_i \overset{\mathbf{L}}{\otimes}_\Lambda T = I_i \otimes_\Lambda T = I_i T$ is a tilting Λ -module of projective dimension at most one.
- (b) $I_i T$ is always a tilting Λ -module of projective dimension at most one, and $\text{End}_\Lambda(I_i T) \simeq \text{End}_\Lambda(T)$.

Proof. (a) Since $\text{Tor}_1^\Lambda(I_i, T) = \text{Tor}_2^\Lambda(S_i, T) = 0$ because $\text{pd } T \leq 1$, we have $I_i \overset{\mathbf{L}}{\otimes}_\Lambda T = I_i \otimes_\Lambda T$. Since we have an exact sequence

$$0 = \text{Tor}_1^\Lambda(S_i, T) \rightarrow I_i \otimes_\Lambda T \rightarrow \Lambda \otimes_\Lambda T \rightarrow S_i \otimes_\Lambda T \rightarrow 0,$$

we have $I_i \otimes_\Lambda T = I_i T$. Thus $I_i \overset{\mathbf{L}}{\otimes}_\Lambda T = I_i T$ is a tilting Λ -module by Lemma II.1.2 and Proposition II.1.4. Since $\text{pd } T \leq 1$ and $\text{pd } T/I_i T \leq 2$ by Proposition II.1.3, we have $\text{pd } I_i T \leq 1$.

(b) By Lemma II.1.1, either $S_i \otimes_\Lambda T = 0$ or $\text{Tor}_1^\Lambda(S_i, T) = 0$ holds. If $S_i \otimes_\Lambda T = 0$, then $I_i T = T$ holds. If $\text{Tor}_1^\Lambda(S_i, T) = 0$, then we apply (a). For the rest we use Lemma II.1.2. $\square$

A left ideal I of Λ is called *cofinite* if $\Lambda/I \in \text{f.l.}\Lambda$, and called *tilting* (respectively, *partial tilting*) if it is a tilting (respectively, partial tilting) Λ -module. Similarly, a *cofinite* (respectively, *(partial) tilting*) *right ideal* of Λ is defined. An ideal I of Λ is called *cofinite tilting* if it is cofinite tilting as a left and right ideal. We denote by $\langle I_1, \dots, I_n \rangle$ the ideal semigroup generated by $I_1, \dots, I_n$. Then we have the following result.

Theorem II.1.6. (a) Any $T \in \langle I_1, \dots, I_n \rangle$ is a cofinite tilting ideal and satisfies $\text{End}_\Lambda(T) = \Lambda$.
 (b) Any cofinite tilting ideal of Λ belongs to $\langle I_1, \dots, I_n \rangle$.
 (c) Any cofinite partial tilting left (respectively, right) ideal of Λ is a cofinite tilting ideal.

Proof. (a) This is a direct consequence of Propositions II.1.4 and II.1.5.

(b)(c) Let T be a cofinite partial tilting left ideal of Λ . If $T \neq \Lambda$, then there exists a simple submodule S_i of Λ/T . Since $\text{Hom}_\Lambda(S_i, \Lambda) = 0$, we have $\text{Ext}_\Lambda^1(S_i, T) \neq 0$. Thus we have $\text{Tor}_1^\Lambda(S_i, T) \simeq D\text{Ext}_\Lambda^1(T, S_i) \simeq \text{Ext}_\Lambda^1(S_i, T) \neq 0$. By Lemma II.1.1, we have $S_i \otimes_\Lambda T = 0$.

Put $U = \mathbf{R}\text{Hom}_\Lambda(I_i, T)$. By Lemma II.1.2, we have that $U \simeq \mathbf{R}\text{Hom}_\Lambda(I_i, \Lambda) \overset{\mathbf{L}}{\otimes}_\Lambda T$ is a partial tilting complex of Λ . Since $\text{pd } I_i \leq 1$ and $\text{Ext}_\Lambda^1(I_i, T) \simeq \text{Ext}_\Lambda^2(S_i, T) \simeq$

$D \operatorname{Hom}_\Lambda(T, S_i) \simeq S_i \otimes_\Lambda T = 0$, we have $U = \operatorname{Hom}_\Lambda(I_i, T)$, which is a partial tilting Λ -module. Since we have a commutative diagram

$$\begin{array}{ccccccc} 0 = \operatorname{Hom}_\Lambda(S_i, \Lambda) & \rightarrow & \Lambda & \rightarrow & \operatorname{Hom}_\Lambda(I_i, \Lambda) & \rightarrow & \operatorname{Ext}_\Lambda^1(S_i, \Lambda) = 0 \\ & & \cup & & \cup & & \\ & & T & \rightarrow & \operatorname{Hom}_\Lambda(I_i, T) & \rightarrow & \operatorname{Ext}_\Lambda^1(S_i, T) \rightarrow 0 \end{array}$$

of exact sequences, U is a cofinite partial tilting left ideal of Λ containing T properly, such that U/T is a direct sum of copies of S_i . By $S_i \otimes_\Lambda T = 0$, we have $T = I_i U$. Thus we have $T \in \langle I_1, \dots, I_n \rangle$ by induction on the length of Λ/T . $\square$

We here pose the following question, where there is a positive answer in the extended Dynkin case [IR].

Question II.1.7. *For any tilting Λ -module T of projective dimension at most one, does there exist some U in $\langle I_1, \dots, I_n \rangle$ such that $\operatorname{add} T = \operatorname{add} U$?*

We have some stronger statements on products of the ideals I_i , generalizing results in the noetherian case from [IR].

Proposition II.1.8. *The following equalities hold for multiplication of ideals.*

- (a) $I_i^2 = I_i$,
- (b) $I_i I_j = I_j I_i$ if there is no arrow between i and j in Q ,
- (c) $I_i I_j I_i = I_j I_i I_j$ if there is precisely one arrow between i and j in Q .

Proof. (a) is obvious.

Parts (b) and (c) are proved in [IR, 6.12] for module-finite 2-CY algebras. Here we give a direct proof for an arbitrary preprojective algebra Λ associated with a finite quiver Q without oriented cycles. Let $I_{i,j} = \Lambda(1 - e_i - e_j)\Lambda$. Then any product of ideals I_i and I_j contains $I_{i,j}$. If there is no arrow from i to j , then $\Lambda/I_{i,j}$ is semisimple. Thus $I_i I_j$ and $I_j I_i$ are contained in $I_{i,j}$, and we have $I_i I_j = I_{i,j} = I_j I_i$.

If there is precisely one arrow from i to j , then $\Lambda/I_{i,j}$ is the preprojective algebra of type A_2 . Hence there are two indecomposable projective $\Lambda/I_{i,j}$ -modules, whose Loewy series are $\begin{pmatrix} i \\ j \end{pmatrix}$ and $\begin{pmatrix} j \\ i \end{pmatrix}$. Thus $I_i I_j I_i$ and $I_j I_i I_j$ are contained in $I_{i,j}$, and we have $I_i I_j I_i = I_{i,j} = I_j I_i I_j$. $\square$

Now let W be the Coxeter group associated to the quiver Q , so W has generators $s_1, \dots, s_n$ with relations $s_i^2 = 1$, $s_i s_j = s_j s_i$ if there is no arrow between i and j in Q , and $s_i s_j s_i = s_j s_i s_j$ if there is a precisely one arrow between i and j in Q .

Theorem II.1.9. *There exists a bijection $W \rightarrow \langle I_1, \dots, I_n \rangle$. It is given by $w \mapsto I_w = I_{i_1} I_{i_2} \dots I_{i_k}$ for any reduced expression $w = s_{i_1} s_{i_2} \dots s_{i_k}$.*

Proof. The corresponding result was proved in [IR] in the noetherian case, using a partial order of tilting modules. Here we use instead properties of Coxeter groups.

We first show that the map is well-defined. Take two reduced expressions $w = s_{i_1} s_{i_2} \dots s_{i_k} = s_{j_1} s_{j_2} \dots s_{j_k}$. By [BB, 3.3.1(ii)], two words $s_{i_1} s_{i_2} \dots s_{i_k}$ and $s_{j_1} s_{j_2} \dots s_{j_k}$ can be connected by a sequence of the following operations: (i) replace $s_i s_j$ by $s_j s_i$ (there is no arrow from i to j), (ii) replace $s_i s_j s_i$ by $s_j s_i s_j$ (there is precisely one arrow from i to j). Consequently, by Proposition II.1.8(b)(c), we have $I_{i_1} I_{i_2} \dots I_{i_k} = I_{j_1} I_{j_2} \dots I_{j_k}$. Thus the map is well-defined.

Next we show that the map is surjective. For any $I \in \langle I_1, \dots, I_n \rangle$, take an expression $I = I_{i_1} I_{i_2} \dots I_{i_k}$ with a minimal number k . Let $w = s_{i_1} s_{i_2} \dots s_{i_k}$. By [BB, 3.3.1(i)], a reduced expression of w is obtained from the word $s_{i_1} s_{i_2} \dots s_{i_k}$ by a sequence of the operations (i)(ii) above and (iii) remove $s_i s_i$. By Proposition II.1.8, the operation (iii) can not appear since k is minimal. Thus $w = s_{i_1} s_{i_2} \dots s_{i_k}$ is a reduced expression, and we have $I = I_w$.

Finally we show that the map is injective in a similar way as in [IR]. Let $\mathcal{E} = \mathbf{K}^b(\text{pr } \Lambda)$. For any i , we have an autoequivalence $I_i \overset{\mathbf{L}}{\otimes}_{\Lambda}$ of $\mathcal{E}$ and an automorphism $[I_i \overset{\mathbf{L}}{\otimes}_{\Lambda}]$ of the Grothendieck group $K_0(\mathcal{E})$. By [IR, proof of 6.6], we have the action $s_i \mapsto [I_i \overset{\mathbf{L}}{\otimes}_{\Lambda}]$ of W on $K_0(\mathcal{E}) \otimes_{\mathbb{Z}} \mathbb{C}$, which is shown to be faithful [BB, 4.2.7].

For any reduced expression $w = s_{i_1} s_{i_2} \dots s_{i_k}$, we have $I_w = I_{i_1} \overset{\mathbf{L}}{\otimes}_{\Lambda} \dots \overset{\mathbf{L}}{\otimes}_{\Lambda} I_{i_k}$, by Proposition II.1.5(a) and the minimality of k . Thus the action of w on $K_0(\mathcal{E}) \otimes_{\mathbb{Z}} \mathbb{C}$ coincides with $[I_w \overset{\mathbf{L}}{\otimes}_{\Lambda}]$. In particular, if $w, w' \in W$ satisfy $I_w = I_{w'}$, then the actions of w and w' on $K_0(\mathcal{E}) \otimes_{\mathbb{Z}} \mathbb{C}$ coincide, so we have $w = w'$ by the faithfulness of the action. $\square$

We denote by $l(w)$ the length of $w \in W$. We say that an infinite expression $s_{i_1} s_{i_2} \dots s_{i_k} \dots$ is *reduced* if the expression $s_{i_1} s_{i_2} \dots s_{i_k}$ is reduced for any k .

Proposition II.1.10. *Let $w \in W$ and $i \in \{1, \dots, n\}$. If $l(ws_i) > l(w)$, then we have $I_w I_i = I_{ws_i} \subsetneq I_w$. If $l(ws_i) < l(w)$, then we have $I_w I_i = I_w \subsetneq I_{ws_i}$.*

Proof. Let $w = s_{i_1} \dots s_{i_k}$ be a reduced expression. If $l(ws_i) > l(w)$, then $ws_i = s_{i_1} \dots s_{i_k} s_i$ is a reduced expression, so the assertion follows from Theorem II.1.9. If $l(ws_i) < l(w)$, then $u = ws_i$ satisfies $l(us_i) > l(u)$, so $I_u I_i = I_{us_i} = I_w \subsetneq I_u$. $\square$

Let $s_{i_1} s_{i_2} \dots s_{i_k} \dots$ be a (finite or infinite) expression such that $i_k \in \{1, \dots, n\}$. Let

$$w_k = s_{i_1} s_{i_2} \dots s_{i_k}, \quad T_k = I_{w_k} = I_{i_1} I_{i_2} \dots I_{i_k} \quad \text{and} \quad \Lambda_k = \Lambda / T_k.$$

We have a descending chain

$$\Lambda = T_0 \supseteq T_1 \supseteq T_2 \supseteq \dots$$

of cofinite tilting ideals of Λ , and a chain

$$\Lambda_1 \leftarrow \Lambda_2 \leftarrow \Lambda_3 \leftarrow \dots$$

of surjective ring homomorphisms. We have the following properties of the chain.

- Proposition II.1.11.** (a) *If $T_{m-1} \neq T_m$, then Λ_m differs from Λ_{m-1} in exactly one indecomposable summand $\Lambda_m e_{i_m}$.*
 (b) *Let $k \leq m$. Then $\Lambda_k e_{i_k}$ is a projective Λ_m -module if and only if $i_k \notin \{i_{k+1}, i_{k+2}, \dots, i_m\}$.*
 (c) *$T_1 \supsetneq T_2 \supsetneq T_3 \supsetneq \dots$ holds if and only if $s_{i_1} s_{i_2} \dots$ is reduced.*

Proof. (a) This follows from $T_m(1 - e_{i_m}) = T_{m-1} I_{i_m}(1 - e_{i_m}) = T_{m-1}(1 - e_{i_m})$.

(b) If $i_k \notin \{i_{k+1}, \dots, i_m\}$, then $\Lambda_k e_{i_k}$ is a summand of Λ_m by (a), so it is a projective Λ_m -module. Otherwise, take the smallest k' with $k < k' \leq m$ satisfying $i_k = i_{k'}$. Then we have $\Lambda_k e_{i_k} = \Lambda_{k'-1} e_{i_k}$ and that $\Lambda_k e_{i_k}$ is a proper factor module of $\Lambda_{k'} e_{i_k}$ by (a). Hence $\Lambda_k e_{i_k}$ is not a projective Λ_m -module.

(c) This follows from Proposition II.1.10. $\square$

Our next goal is to show that $\text{Ext}_{\Lambda}^1(T_k, T_m) = 0$ for $k \leq m$. For this the following result will be useful.

Lemma II.1.12. *Let the notation and assumptions be as above. Then ${}^{\perp > 0} T_{m-1} \subseteq {}^{\perp > 0} T_m$, where ${}^{\perp > 0} T = \{X \in \text{mod } \Lambda \mid \text{Ext}_{\Lambda}^i(X, T) = 0 \text{ for all } i > 0\}$.*

Proof. We can assume $T_{m-1} \neq T_m$. Then we have that $T_{m-1} \otimes_{\Lambda} S_{i_m} \neq 0$. Hence $\text{Tor}_1^{\Lambda}(T_{m-1}, S_{i_m}) = 0$ by Lemma II.1.1, so $T_{m-1} \otimes_{\Lambda} I_{i_m} = T_{m-1} I_{i_m} = T_m$ by Proposition II.1.5. Let $0 \rightarrow P_1 \rightarrow P_0 \rightarrow I_{i_m} \rightarrow 0$ be a projective resolution. We have $\text{Tor}_1^{\Lambda}(T_{m-1}, I_{i_m}) \simeq \text{Tor}_2^{\Lambda}(T_{m-1}, S_{i_m}) = 0$. Applying $T_{m-1} \otimes_{\Lambda}$, we have an exact sequence $0 \rightarrow T_{m-1} \otimes_{\Lambda} P_1 \rightarrow T_{m-1} \otimes_{\Lambda} P_0 \rightarrow T_m \rightarrow 0$. This immediately implies ${}^{\perp > 0} T_{m-1} \subseteq {}^{\perp > 0} T_m$. $\square$

We now have the following consequence.

Proposition II.1.13. *With the above notation and assumptions, we have $\text{Ext}_\Lambda^1(T_k, T_m) = 0$ for $k \leq m$.*

Proof. By Lemma II.1.12 we have ${}^{\perp > 0} T_k \subseteq {}^{\perp > 0} T_{k+1} \subseteq \dots \subseteq {}^{\perp > 0} T_m$. Since T_k is in ${}^{\perp > 0} T_k$, we then have that T_k is in ${}^{\perp > 0} T_m$, and hence $\text{Ext}_\Lambda^1(T_k, T_m) = 0$ for $k \leq m$. $\square$

Later we shall use the following observation.

Lemma II.1.14. *Assume that the expression $s_{i_1} s_{i_2} \dots$ is reduced. Let $T_{k,m} = I_{i_k} \dots I_{i_m}$ if $k \leq m$ and $T_{k,m} = \Lambda$ otherwise. Then we have $\text{Hom}_\Lambda(T_k, T_m) \simeq T_{k+1,m} = \{x \in \Lambda \mid T_k x \subseteq T_m\}$ and $\text{Hom}_\Lambda(\Lambda_k, \Lambda_m) \simeq T_{k+1,m}/T_m$.*

Proof. Let $U = \{x \in \Lambda \mid T_k x \subseteq T_m\} \supseteq T_{k+1,m}$.

If $k \geq m$, then clearly $U = \Lambda = T_{k+1,m}$ holds, and $\text{Hom}_\Lambda(T_k, T_m) \subseteq \text{End}_\Lambda(T_k) \simeq \Lambda$ by Theorem II.1.6. Thus $\text{Hom}_\Lambda(T_k, T_m) \simeq \Lambda$.

We assume $k < m$. Since $T_m = T_k \overset{\mathbf{L}}{\otimes}_\Lambda T_{k+1,m}$ holds by Proposition II.1.5(a) and Lemma II.1.2, we have $\mathbf{R}\text{Hom}_\Lambda(T_k, T_m) = \mathbf{R}\text{Hom}_\Lambda(T_k, (T_k \overset{\mathbf{L}}{\otimes}_\Lambda T_{k+1,m})) = \mathbf{R}\text{Hom}_\Lambda(T_k, T_k) \overset{\mathbf{L}}{\otimes}_\Lambda T_{k+1,m} = \Lambda \overset{\mathbf{L}}{\otimes}_\Lambda T_{k+1,m} = T_{k+1,m}$. In particular, we have $\text{Hom}_\Lambda(T_k, T_m) = T_{k+1,m}$. On the other hand, we have a commutative diagram

$$\begin{array}{ccc} \Lambda & \rightarrow & \text{Hom}_\Lambda(T_k, \Lambda) \\ \cup & & \cup \\ U & \rightarrow & \text{Hom}_\Lambda(T_k, T_m) \simeq T_{k+1,m}, \end{array}$$

where the horizontal map is given by $x \mapsto (\cdot x)$ for any $x \in \Lambda$, which is injective. Thus we have $U \subseteq T_{k+1,m}$, and so $U = T_{k+1,m}$.

Now we show the second equality. For any $f \in \text{Hom}_\Lambda(\Lambda_k, \Lambda_m)$, there exists a unique element $x \in \Lambda_m$ such that $f(y) = yx$ for any $y \in \Lambda$. Since $T_k x \subseteq T_m$ holds, we have $x \in U$. Thus we have $\text{Hom}_\Lambda(\Lambda_k, \Lambda_m) \simeq U/T_m = T_{k+1,m}/T_m$. $\square$

II.2. Cluster tilting objects for preprojective algebras.

Let again Λ be the completion of the preprojective algebra of a finite connected non-Dynkin quiver without loops over the field k . We show that for a large class of cofinite tilting ideals I in Λ we have that Λ/I is a finite dimensional k -algebra which is Gorenstein of dimension at most one, and the categories $\text{Sub } \Lambda/I$ and $\underline{\text{Sub}} \Lambda/I$ are stably 2-CY and 2-CY respectively. We describe some cluster tilting objects in these categories, using tilting ideals. We also describe cluster tilting subcategories in the derived 2-CY abelian category $\text{f.l. } \Lambda$, which have an infinite number of nonisomorphic indecomposable objects. Hence we get examples of cluster structures with infinite clusters (see [KR2] for other examples).

We start with investigating Λ/T for our special cofinite tilting ideals T as a module over Λ and over the factor ring Λ/U for a cofinite tilting ideal U contained in T .

Lemma II.2.1. *Let T and $U = TU'$ be cofinite tilting ideals in Λ . Then $\text{Ext}_\Lambda^1(\Lambda/T, \Lambda/U) = 0 = \text{Ext}_\Lambda^1(\Lambda/U, \Lambda/T)$.*

Proof. Consider the exact sequence $0 \rightarrow U \rightarrow \Lambda \rightarrow \Lambda/U \rightarrow 0$. Applying $\text{Hom}_\Lambda(\Lambda/T, \cdot)$, we have an exact sequence

$$\text{Ext}_\Lambda^1(\Lambda/T, \Lambda) \rightarrow \text{Ext}_\Lambda^1(\Lambda/T, \Lambda/U) \rightarrow \text{Ext}_\Lambda^2(\Lambda/T, U).$$

We have $\text{Ext}_\Lambda^1(\Lambda/T, \Lambda) = 0$ by Proposition II.1.3. It follows from Corollary II.1.13 that $\text{Ext}_\Lambda^1(T, U) = 0$. Since $\text{Ext}_\Lambda^2(\Lambda/T, U) \simeq \text{Ext}_\Lambda^1(T, U) = 0$, it follows that $\text{Ext}_\Lambda^1(\Lambda/T, \Lambda/U) = 0$. Since Λ is derived 2-CY it follows that also $\text{Ext}_\Lambda^1(\Lambda/U, \Lambda/T) = 0$. $\square$

Using this lemma we obtain more information on Λ/T .

- Proposition II.2.2.** (a) For a cofinite ideal T in Λ with $\text{Ext}_\Lambda^1(\Lambda/T, \Lambda/T) = 0$, the algebra Λ/T is Gorenstein of dimension at most one.
 (b) For a cofinite tilting ideal T in Λ , the factor algebra Λ/T is Gorenstein of dimension at most one.

Proof. (a) Consider the exact sequence $0 \rightarrow \Omega_\Lambda(D(\Lambda/T)) \rightarrow P \rightarrow D(\Lambda/T) \rightarrow 0$ with a projective Λ -module P . Using Lemma II.2.1 and [CE], we have $\text{Tor}_1^\Lambda(\Lambda/T, D(\Lambda/T)) \simeq D \text{Ext}_{\Lambda^{\text{op}}}^1(\Lambda/T, \Lambda/T) = 0$. Applying $\Lambda/T \otimes_\Lambda$ to the above exact sequence, we get the exact sequence $0 \rightarrow \Lambda/T \otimes_\Lambda \Omega_\Lambda(D(\Lambda/T)) \rightarrow \Lambda/T \otimes_\Lambda P \rightarrow \Lambda/T \otimes_\Lambda D(\Lambda/T) \rightarrow 0$. The Λ/T -module $\Lambda/T \otimes_\Lambda P$ is projective. To see that also $\Lambda/T \otimes_\Lambda \Omega_\Lambda(D(\Lambda/T))$ is a projective Λ/T -module, we show that the functor $\text{Hom}_{\Lambda/T}(\Lambda/T \otimes_\Lambda \Omega_\Lambda(D(\Lambda/T)), _) \simeq \text{Hom}_\Lambda(\Omega_\Lambda(D(\Lambda/T)), _)$ is exact on $\text{mod } \Lambda/T$. This follows from the functorial isomorphisms

$$\begin{aligned} \text{Ext}_\Lambda^1(\Omega_\Lambda(D(\Lambda/T)), _) &\simeq \text{Ext}_\Lambda^2(D(\Lambda/T), _) \\ &\simeq D \text{Hom}_\Lambda(_, D(\Lambda/T)) \\ &\simeq D \text{Hom}_{\Lambda/T}(_, D(\Lambda/T)) \simeq \text{id}_{\text{mod } \Lambda/T} \end{aligned}$$

Hence we conclude that $\text{pd}_{\Lambda/T} D(\Lambda/T) \leq 1$. Then it is well known and easy to see that $\text{pd}_{(\Lambda/T)^{\text{op}}} D(\Lambda/T) \leq 1$, so that by definition Λ/T is Gorenstein of dimension at most one.

(b) This is a direct consequence of (a) and Lemma II.2.1. $\square$

When Λ/T is Gorenstein of dimension at most one, the category of Cohen-Macaulay modules is the category $\text{Sub}(\Lambda/T)$ of first syzygy modules (see [AR1, H2]). It is known that $\text{Sub}(\Lambda/T)$ is a Frobenius category, with $\text{add}(\Lambda/T)$ being the category of projective and injective objects, and the stable category $\underline{\text{Sub}}(\Lambda/T)$ is triangulated [H1]. Moreover $\text{Sub}(\Lambda/T)$ is an extension closed subcategory of $\text{mod } \Lambda/T$ by Corollary I.2.8, since $\text{id}_{\Lambda/T} \Lambda/T \leq 1$ and $\text{Ext}_{\Lambda/T}^1(\Lambda/T, \Lambda/T) = 0$. But to show that the stably 2-CY property is deduced from f.l. Λ being derived 2-CY we need that $\text{Sub } \Lambda/T$ is also extension closed in f.l. Λ .

Proposition II.2.3. Let T be a cofinite ideal with $\text{Ext}_\Lambda^1(\Lambda/T, \Lambda/T) = 0$ (for example a cofinite tilting ideal).

- (a) $\text{Ext}_\Lambda^1(\Lambda/T, X) = 0 = \text{Ext}_\Lambda^1(X, \Lambda/T)$ for all X in $\text{Sub } \Lambda/T$.
 (b) $\text{Sub } \Lambda/T$ is an extension closed subcategory of f.l. Λ .
 (c) $\text{Sub } \Lambda/T$ and $\underline{\text{Sub}} \Lambda/T$ are stably 2-CY and 2-CY respectively.

Proof. (a) For X in $\text{Sub } \Lambda/T$ we have an exact sequence $0 \rightarrow X \rightarrow P \rightarrow Y \rightarrow 0$ with Y in $\text{Sub } \Lambda/T$ and P in $\text{add } \Lambda/T$. Applying $\text{Hom}_\Lambda(\Lambda/T, _)$ $\simeq \text{Hom}_{\Lambda/T}(\Lambda/T, _)$, the sequence does not change. Since $\text{Ext}_\Lambda^1(\Lambda/T, \Lambda/T) = 0$, we conclude that $\text{Ext}_\Lambda^1(\Lambda/T, X) = 0$. Hence $\text{Ext}_\Lambda^1(X, \Lambda/T) = 0$ by the derived 2-CY property of f.l. Λ .

(b) Let $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$ be an exact sequence in f.l. Λ , with X and Z in $\text{Sub } \Lambda/T$. Then we have a monomorphism $X \rightarrow P$, with P in $\text{add } \Lambda/T$. Since $\text{Ext}_\Lambda^1(Z, P) = 0$ by (a), we have a commutative diagram of exact sequences

$$\begin{array}{ccccccc} 0 & \rightarrow & X & \rightarrow & Y & \rightarrow & Z \rightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \rightarrow & P & \rightarrow & P \oplus Z & \rightarrow & Z \rightarrow 0 \end{array}$$

Thus Y is a submodule of $P \oplus Z \in \text{Sub } \Lambda/T$, and we have $Y \in \text{Sub } \Lambda/T$.

(c) Since $\text{Sub } \Lambda/T$ is extension closed in f.l. Λ , we have $\text{Ext}_{\text{Sub } \Lambda/T}^1(X, Y) = \text{Ext}_\Lambda^1(X, Y)$. Since $\text{Sub } \Lambda/T$ is Frobenius, it follows from Proposition I.1.1, that $\text{Sub } \Lambda/T$ is stably 2-CY, since f.l. Λ is derived 2-CY, and so $\underline{\text{Sub}} \Lambda/T$ is 2-CY. $\square$

We now want to investigate the cluster tilting objects in $\text{Sub } \Lambda/T$ and $\underline{\text{Sub}} \Lambda/T$ for certain tilting ideals T , and later also the cluster tilting subcategories of f.l. Λ . The following observation will be useful.

Lemma II.2.4. *Let Δ be a finite dimensional algebra and M a Δ -module which is a generator. Let $\Gamma = \text{End}_\Delta(M)$, and assume $\text{gl.dim } \Gamma \leq 3$ and $\text{pd}_\Gamma D(M) \leq 1$. Then for any X in $\text{mod } \Delta$ there is an exact sequence $0 \rightarrow M_1 \rightarrow M_0 \rightarrow X \rightarrow 0$, with M_0 and M_1 in $\text{add } M$.*

Proof. Let X be in $\text{mod } \Delta$, and consider the exact sequence $0 \rightarrow X \rightarrow I_0 \rightarrow I_1$ where I_0 and I_1 are injective. Apply $\text{Hom}_\Delta(M, _)$ to get an exact sequence $0 \rightarrow \text{Hom}_\Delta(M, X) \rightarrow \text{Hom}_\Delta(M, I_0) \rightarrow \text{Hom}_\Delta(M, I_1)$. Since by assumption $\text{pd}_\Gamma \text{Hom}_\Delta(M, I_i) \leq 1$ for $i = 0, 1$ and $\text{gl.dim } \Gamma \leq 3$, we obtain $\text{pd}_\Gamma \text{Hom}_\Delta(M, X) \leq 1$. Hence we have an exact sequence $0 \rightarrow P_1 \rightarrow P_0 \rightarrow \text{Hom}_\Delta(M, X) \rightarrow 0$ in $\text{mod } \Gamma$ with P_0 and P_1 projective. This sequence is the image under the functor $\text{Hom}_\Delta(M, _)$ of the complex $0 \rightarrow M_1 \rightarrow M_0 \rightarrow X \rightarrow 0$ in $\text{mod } \Delta$, with M_0 and M_1 in $\text{add } M$. Since M is assumed to be a generator, this complex must be exact, and we have our desired exact sequence. $\square$

Let now $\Lambda = T_0 \supsetneq T_1 \supsetneq T_2 \supsetneq \cdots$ be a strict descending chain of tilting ideals corresponding to a (finite or infinite) reduced expression $s_{i_1} s_{i_2} s_{i_3} \cdots$. We want to describe some natural cluster tilting objects for the algebras $\Lambda_m = \Lambda/T_m$. Let

$$\Lambda_k = \Lambda/T_k \text{ and } M_m = \bigoplus_{k=0}^m \Lambda_k,$$

and $\Gamma = \text{End}_{\Lambda_m}(M_m)$. The following will be essential.

Proposition II.2.5. *With the above notation we have the following.*

- (a) *For X in $\text{mod } \Lambda_m$ there is an exact sequence $0 \rightarrow N_1 \rightarrow N_0 \rightarrow X \rightarrow 0$ in $\text{mod } \Lambda_m$, with N_i in $\text{add } M_m$ for $i = 1, 2$.*
- (b) *$\text{gl.dim } \Gamma \leq 3$.*

Proof. We prove (a) and (b) by induction on m . Assume first that $m = 1$. Then $\Lambda_1 = \Lambda/T_1$, which is a simple Λ_1 -module. Since $M_1 = \Lambda/T_1$, (a) and (b) are trivially satisfied.

Assume now that $m > 1$ and that (a) and (b) have been proved for $m - 1$. Then we first prove (b) for m . Note that since there are no loops for Λ , we have $T_{m-1}J \subseteq T_m$ where J is the Jacobson radical of Λ , so that $J\Lambda_m$ is a Λ_{m-1} -module (*).

For an indecomposable object X in $\mathcal{M}_m = \text{add } M_m$, let $f: C_0 \rightarrow X$ be a minimal right almost split map in $\mathcal{M}_m$. We first assume that X is not a projective Λ_m -module. Then f must be surjective. An indecomposable object which is in $\mathcal{M}_m$ but not in $\mathcal{M}_{m-1}$ is a projective Λ_m -module, so we can write $C_0 = C'_0 \oplus P$ where $C'_0 \in \mathcal{M}_{m-1}$ and P is a projective Λ_m -module. Since f is right minimal, we have $\text{Ker } f \subseteq C'_0 \oplus JP$, so that $\text{Ker } f$ is a Λ_{m-1} -module by (*). It follows by the induction assumption that there is an exact sequence $0 \rightarrow C_2 \rightarrow C_1 \rightarrow \text{Ker } f \rightarrow 0$ with C_1 and C_2 in $\mathcal{M}_{m-1}$. Hence we have an exact sequence $0 \rightarrow C_2 \rightarrow C_1 \rightarrow C_0 \rightarrow X \rightarrow 0$. Applying $\text{Hom}_\Lambda(M_m, _)$ gives an exact sequence

$$0 \rightarrow \text{Hom}_\Lambda(M_m, C_2) \rightarrow \text{Hom}_\Lambda(M_m, C_1) \rightarrow \text{Hom}_\Lambda(M_m, C_0) \rightarrow \text{Hom}_\Lambda(M_m, X) \rightarrow S \rightarrow 0.$$

Then the module S , which is a simple module in the top of $\text{Hom}_\Lambda(M_m, X)$ in $\text{mod } \Gamma$, has projective dimension at most 3.

Assume now that X is a projective Λ_m -module. Then by (*) we have that JX is in $\text{mod } \Lambda_{m-1}$. By the induction assumption there is then an exact sequence $0 \rightarrow C_1 \rightarrow C_0 \rightarrow JX \rightarrow 0$, with C_0 and C_1 in $\mathcal{M}_{m-1}$. Hence we have an exact sequence $0 \rightarrow C_1 \rightarrow C_0 \rightarrow X$. Applying $\text{Hom}_\Lambda(M_m, _)$ gives the exact sequence

$$0 \rightarrow \text{Hom}_\Lambda(M_m, C_1) \rightarrow \text{Hom}_\Lambda(M_m, C_0) \rightarrow \text{Hom}_\Lambda(M_m, X) \rightarrow S \rightarrow 0$$

where S is the simple top of $\text{Hom}_\Lambda(M_m, X)$, and hence $\text{pd}_\Gamma S \leq 2$. It now follows that $\text{gl.dim } \Gamma_m \leq 3$.

We now want to show (a) for m . By Proposition II.2.2 we have an exact sequence $0 \rightarrow P_1 \rightarrow P_0 \rightarrow D(\Lambda_m) \rightarrow 0$ in $\text{mod } \Lambda_m$, where P_0 and P_1 are projective Λ_m -modules. By Lemma II.2.1 we have $\text{Ext}_\Lambda^1(M_m, \Lambda_m) = 0$. Applying $\text{Hom}_\Lambda(M_m, _)$ gives the exact sequence

$$0 \rightarrow \text{Hom}_\Lambda(M_m, P_1) \rightarrow \text{Hom}_\Lambda(M_m, P_0) \rightarrow \text{Hom}_\Lambda(M_m, D(\Lambda_m)) \rightarrow 0$$

Since $\text{Hom}_\Lambda(M_m, D(\Lambda_m)) \simeq D(M_m)$, we have $\text{pd}_{\Gamma_m} D(M_m) \leq 1$. Now our desired result follows from Lemma II.2.4. $\square$

We can now describe some cluster tilting objects in $\text{Sub } \Lambda_m$ and $\underline{\text{Sub}} \Lambda_m$.

Theorem II.2.6. *With the above notation, M_m is a cluster tilting object in $\text{Sub } \Lambda_m$ and in $\underline{\text{Sub}} \Lambda_m$.*

Proof. We already have that $\text{Ext}_\Lambda^1(M_m, M_m) = 0$ by Lemma II.2.1, so $\text{Ext}_{\Lambda_m}^1(M_m, M_m) = 0$. Note that $\text{Sub } \Lambda_m = \{X \in \text{mod } \Lambda_m \mid \text{Ext}_{\Lambda_m}^1(X, \Lambda_m) = 0\}$ because Λ_m is a cotilting module with $\text{id } \Lambda_m \leq 1$. Since Λ_m is a summand of M_m , we have that M_m is in $\text{Sub } \Lambda_m$. Assume then that $\text{Ext}_{\Lambda_m}^1(X, M_m) = 0$ for X in $\text{mod } \Lambda_m$. By Proposition II.2.5(a) there is an exact sequence $0 \rightarrow C_1 \rightarrow C_0 \rightarrow X \rightarrow 0$ with C_1 and C_0 in $\text{add } M_m$, which must split by our assumption. Hence X is in $\text{add } M_m$, and it follows that M_m is a cluster tilting object in $\text{Sub } \Lambda_m$. It then follows as usual that it is a cluster tilting object also in $\underline{\text{Sub}} \Lambda_m$. $\square$

We have now obtained a large class of 2-CY categories $\text{Sub } \Lambda/I_w$ and $\underline{\text{Sub}} \Lambda/I_w$ defined via elements w of the associated Coxeter group W , along with cluster tilting objects associated with reduced expressions of elements in W . We call these *standard cluster tilting objects* for $\text{Sub } \Lambda/I_w$ or $\underline{\text{Sub}} \Lambda/I_w$. We can also describe cluster tilting subcategories with an infinite number of nonisomorphic indecomposable objects in the categories f.l. Λ .

Theorem II.2.7. *With the above notation, assume that each i occurs an infinite number of times in $i_1, i_2, \dots$. Then $\mathcal{M} = \text{add}\{\Lambda_m \mid 0 \leq m\}$ is a cluster tilting subcategory of f.l. Λ .*

Proof. We already know that $\text{Ext}_\Lambda^1(\Lambda_k, \Lambda_m) = 0$ for all k and m . Let now X be indecomposable in f.l. Λ . Then X is a Λ/J^k -module for some k . We have $J = I_1 \cap \dots \cap I_n \supseteq I_1 \cdots I_n$, where $1, \dots, n$ are the vertices in the quiver. By our assumptions we have $J^k \supseteq T_m$ for some m , so that X is a Λ_m -module. Consider the exact sequence $0 \rightarrow C_1 \rightarrow C_0 \rightarrow X \rightarrow 0$ in $\text{mod } \Lambda_m$, with C_1 and C_0 in $\text{add } M_m$, obtained from Proposition II.2.5. Assume that $\text{Ext}_\Lambda^1(X, \mathcal{M}) = 0$. Since also $\text{Ext}_{\Lambda_m}^1(X, \mathcal{M}_m) = 0$, the sequence splits, so that X is in $\mathcal{M}_m$ and hence in $\mathcal{M}$.

It only remains to show that $\mathcal{M}$ is functorially finite. So let X be in f.l. Λ . Using the above exact sequence $0 \rightarrow C_1 \rightarrow C_0 \rightarrow X \rightarrow 0$, we get the exact sequence

$$0 \rightarrow (C, C_1) \rightarrow (C, C_0) \rightarrow (C, X) \rightarrow \text{Ext}_\Lambda^1(C, C_1) = 0$$

for C in $\mathcal{M}$. Hence $\mathcal{M}$ is contravariantly finite.

For X in f.l. Λ , take the left $(\text{Sub } \mathcal{M})$ -approximation $X \rightarrow Y$ and choose m such that $Y \in \text{Sub } \Lambda_m$. For any $Y \in \text{Sub } \Lambda_m$, there exists an exact sequence $0 \rightarrow Y \rightarrow C_0 \rightarrow C_1 \rightarrow 0$ with $C_i \in \mathcal{M}_m$ by Proposition I.1.7(a). Then the composition $X \rightarrow Y \rightarrow C_0$ is a left $\mathcal{M}$ -approximation since $\text{Ext}_\Lambda^1(C_1, \mathcal{M}) = 0$, and hence $\mathcal{M}$ is also covariantly finite. $\square$

Summarizing our results, we have the following.

Theorem II.2.8. (a) *For any $w \in W$, we have a stably 2-CY category $\mathcal{C}_w = \text{Sub } \Lambda/I_w$.*
 (b) *For any reduced expression $w = s_{i_1} \cdots s_{i_m}$ of $w \in W$, we have a cluster tilting object $\bigoplus_{k=1}^m \Lambda/I_{s_{i_1} \cdots s_{i_k}}$ in $\mathcal{C}_w$. In particular the number of non-isomorphic indecomposable summands in any cluster tilting object is $l(w)$.*
 (c) *For any infinite reduced expression $s_{i_1} s_{i_2} \cdots$ such that each i occurs an infinite number of times in $i_1, i_2, \dots$, we have a cluster tilting subcategory $\text{add}\{\Lambda/I_{s_{i_1} \cdots s_{i_k}} \mid 0 \leq k\}$ in f.l. Λ .*

We end this section by showing that the subcategories $\text{Sub } \Lambda/I_w$ can be characterized using torsionfree classes.

Theorem II.2.9. *Let Λ be the completed preprojective algebra of a connected non-Dynkin quiver without loops. Let $\mathcal{C}$ be a torsionfree class in f.l. Λ with some cluster tilting object. Then we have $\mathcal{C} = \text{Sub } \Lambda/I_w$ for some element w in the Coxeter group associated with Λ .*

Proof. We first prove that if M is a cluster tilting object in $\mathcal{C}$, then $\mathcal{C} = \text{Sub } M$. We only have to show $\mathcal{C} \subset \text{Sub } M$. For any $X \in \mathcal{C}$, take a projective resolution $\text{Hom}_\Lambda(M, N) \rightarrow \text{Ext}_\Lambda^1(M, X) \rightarrow 0$ (*) of $\text{End}_\Lambda(M)$ -modules with $N \in \text{add } M$. Replacing M in (*) by N , we get an exact sequence $0 \rightarrow X \rightarrow Y \rightarrow N \rightarrow 0$ as the image of the identity $1_N \in \text{End}_\Lambda(N)$ in $\text{Ext}_\Lambda^1(N, X)$. Since $\mathcal{C}$ is extension closed, we have $Y \in \mathcal{C}$. We have an exact sequence $\text{Hom}_\Lambda(M, N) \rightarrow \text{Ext}_\Lambda^1(M, X) \rightarrow \text{Ext}_\Lambda^1(M, Y) \rightarrow \text{Ext}_\Lambda^1(M, N) = 0$. Since (*) is exact, we have $\text{Ext}_\Lambda^1(M, Y) = 0$. Thus we have $Y \in \text{add } M$ and $X \in \text{Sub } M$.

Let now I be the annihilator $\text{ann}_\Lambda M$ of M in Λ . Then I is clearly a cofinite ideal in Λ , and $\text{ann}_{\Lambda/I} M = 0$. Further $\text{Sub } M$ is extension closed also in $\text{mod } \Lambda/I$. Hence the direct sum A of one copy of each of the non-isomorphic indecomposable Ext-injective Λ/I -modules in $\text{Sub } M$ is a cotilting Λ/I -module satisfying $\text{id}_{\Lambda/I} A \leq 1$ and $\text{Sub } M = \text{Sub } A$ by [Sm]. Since $\text{Sub } M$ is extension closed in the derived 2-CY category f.l. Λ , the Ext-injective Λ/I -modules in $\text{Sub } M$ coincide with the Ext-projective ones, which are the projective Λ/I -modules. Hence we have that A is a progenerator of Λ/I and $\text{Sub } M = \text{Sub } \Lambda/I$. Since $\text{Sub } \Lambda/I$ is extension closed in f.l. Λ , we have $\text{Ext}_\Lambda^1(\Lambda/I, \Lambda/I) = 0$.

By Theorem II.1.6, we only have to show that I is a partial tilting left ideal. By Bongartz completion, we only have to show $\text{Ext}_\Lambda^1(I, I) = 0$. The natural surjection $\Lambda \rightarrow \Lambda/I$ clearly induces a surjection $\text{Hom}_\Lambda(\Lambda/I, \Lambda/I) \rightarrow \text{Hom}_\Lambda(\Lambda, \Lambda/I)$. Since Λ is derived 2-CY, we have injections $\text{Ext}_\Lambda^2(\Lambda/I, \Lambda) \rightarrow \text{Ext}_\Lambda^2(\Lambda/I, \Lambda/I)$ and $\text{Ext}_\Lambda^1(I, \Lambda) \rightarrow \text{Ext}_\Lambda^1(I, \Lambda/I)$. Using the exact sequence $0 \rightarrow I \rightarrow \Lambda \rightarrow \Lambda/I \rightarrow 0$, we have a commutative diagram

$$\begin{array}{ccccccc}
 & & & \text{Ext}_\Lambda^1(\Lambda/I, \Lambda/I) = 0 & & & \\
 & & & \uparrow & & & \\
 \text{Hom}_\Lambda(I, \Lambda) & \rightarrow & \text{Hom}_\Lambda(I, \Lambda/I) & \rightarrow & \text{Ext}_\Lambda^1(I, I) & \rightarrow & \text{Ext}_\Lambda^1(I, \Lambda) \rightarrow \text{Ext}_\Lambda^1(I, \Lambda/I) \\
 \uparrow & & \uparrow & & \uparrow & & \\
 \Lambda & \rightarrow & \Lambda/I & \rightarrow & 0 & &
 \end{array}$$

of exact sequences. Thus we have $\text{Ext}_\Lambda^1(I, I) = 0$. $\square$

Note that we have proved that an extension closed subcategory of f.l. Λ of the form $\text{Sub } X$ for some X in f.l. Λ with some cluster tilting object must be $\text{Sub } \Lambda/I_w$ for some element w in the Coxeter group associated with Λ .

We point out that there are other extension closed subcategories of f.l. Λ with some cluster tilting object. Let Q be an extended Dynkin quiver and Q' a Dynkin subquiver, and Λ and Λ' the corresponding completed preprojective algebras. Then clearly $\text{mod } \Lambda' = \text{f.l. } \Lambda'$ is an extension closed subcategory of f.l. Λ . Hence any extension closed subcategory of $\text{mod } \Lambda'$ is extension closed in f.l. Λ , so Example 1 in Section I.3 is an example of an extension closed stably 2-CY subcategory of f.l. Λ with some cluster tilting object, but which is not closed under submodules.

II.3. Realization of cluster categories and stable categories for preprojective algebras of Dynkin type.

In this section we show that for an appropriate choice of T as a product of tilting ideals $I_j = \Lambda(1 - e_j)\Lambda$, any cluster category is equivalent to some $\text{Sub } \Lambda/T$. In particular, any cluster category can be realized as the stable category of a Frobenius category with finite dimensional homomorphism spaces. We also show that the stable categories for preprojective algebras of Dynkin type can be realized this way.

Let Q be a finite connected quiver without loops, kQ the associated path algebra, and Λ the completion of the preprojective algebra of Q . Choose a complete set of orthogonal primitive idempotents $e_1, \dots, e_n$ of kQ . We can assume that $e_i(kQ)e_j = 0$ for any $i > j$. We regard $e_1, \dots, e_n$ as a complete set of orthogonal primitive idempotents of Λ , and let as before $I_i = \Lambda(1 - e_i)\Lambda$.

Assume first that Q is not Dynkin. We consider an exact stably 2-CY category associated to the square w^2 of a Coxeter element $w = s_1 s_2 \cdots s_n \in W$. Let $\Lambda_i = \Lambda / I_1 I_2 \cdots I_i$ and $\Lambda_{i+n} = \Lambda / I_1 I_2 \cdots I_n I_1 \cdots I_i$ for $1 \leq i \leq n$. We have seen in Section II.2 that $\text{Sub } \Lambda_{2n}$, and also $\underline{\text{Sub}} \Lambda_{2n}$, has a cluster tilting object $M = \bigoplus_{i=1}^{2n} \Lambda_i$.

We shall need the following.

Lemma II.3.1. *Assume that Q is not Dynkin. Then $I_1 \cdots I_n I_1 \cdots I_n I_1 \cdots$ gives rise to a strict descending chain of tilting ideals. In particular, $s_1 \cdots s_n s_1 \cdots s_n s_1 \cdots$ is reduced.*

Proof. Assume to the contrary that the descending chain of ideals is not strict. Let $T_i = I_1 \cdots I_i$ and $U_i = I_1 \cdots I_{i-1} I_{i+1} \cdots I_n$ for $i = 1, \dots, n$. Then we have $T_n^k T_{i-1} = T_n^k T_i$ for some $i = 1, \dots, n$ and $k \geq 0$, where $T_0 = \Lambda$. Hence we obtain $T_n^{k+1} = T_n^k U_i$. Then we get $T_n^{k+m} = T_n^k U_i^m$ for any $m > 0$. Since $U_i e_i = \Lambda e_i$ and $J \supseteq T_n$, we have $J^{m+k} e_i \supseteq T_n^{m+k} e_i = T_n^k U_i^m e_i = T_n^k e_i$. Since $(\Lambda / T_n^k) e_i$ has finite length, we have $J^{m+k} e_i = T_n^k e_i$ for m sufficiently large. Thus we have $T_n^k e_i = 0$, which is a contradiction since Λe_i has infinite length. We have the latter assertion from Proposition II.1.11. $\square$

We have the following.

Proposition II.3.2. *Let Q be a finite connected non-Dynkin quiver without oriented cycles and with vertices $1, \dots, n$ ordered as above. Let $\Lambda_{2n} = \Lambda / (I_1 \cdots I_n)^2$. Then $\Lambda_n = \Lambda / I_1 \cdots I_n$ is a cluster tilting object in $\underline{\text{Sub}} \Lambda_{2n}$ with $\text{End}_{\underline{\text{Sub}} \Lambda_{2n}}(\Lambda_n) \simeq kQ$.*

Proof. Since the associated chain of ideals is strict descending by Lemma II.3.1, our general theory applies. We have a cluster tilting object $\bigoplus_{i=1}^{2n} \Lambda_i$ in $\underline{\text{Sub}} \Lambda_{2n}$ by Theorem II.2.6. We have $\text{add } \bigoplus_{i=1}^{2n} \Lambda_i = \Lambda_n \oplus \Lambda_{2n}$ in f.l. Λ by Proposition II.1.11. Thus Λ_n is a cluster tilting object in $\underline{\text{Sub}} \Lambda_{2n}$.

Note that the path algebra kQ is, in a natural way, a factor algebra of Λ , and kQ is hence a Λ -module. We want to show that the Λ -modules Λ_n and kQ are isomorphic.

Let P_j be the indecomposable projective Λ -module corresponding to the vertex j . Then $I_{j+1} \cdots I_n P_j = P_j$ and $I_j P_j = J P_j$, the smallest submodule of P_j such that the corresponding factor has only composition factors S_j . Further, $I_{j-1} I_j P_j$ is the smallest submodule of $I_j P_j = J P_j$ such that the factor has only composition factors S_{j-1} , etc. By our choice of ordering, this means that the paths starting at j , with decreasing indexing on the vertices, give a basis for $P_j / I_1 \cdots I_n P_j$. In other words, we have $P_j / I_1 \cdots I_n P_j \simeq (kQ) e_j$. Hence the Λ -modules $\Lambda_n = \Lambda / I_1 \cdots I_n$ and kQ are isomorphic, so that $\text{End}_{\Lambda}(\Lambda_n) \simeq kQ$.

It remains to show $\text{End}_{\Lambda_{2n}}(\Lambda_n) \simeq \text{End}_{\underline{\text{Sub}} \Lambda_{2n}}(\Lambda_n)$. By Lemma II.1.14, any morphism from Λ_n to Λ_{2n} is given by a right multiplication of an element in $(I_1 \cdots I_n) / (I_1 \cdots I_n)^2$. This implies $\text{Hom}_{\Lambda}(\Lambda_n, \Lambda_{2n}) \text{Hom}_{\Lambda}(\Lambda_{2n}, \Lambda_n) = 0$. Thus we have the assertion. $\square$

We now show that we have the same kind of result for Dynkin quivers.

Proposition II.3.3. *Let Q' be a Dynkin quiver with vertices $1, \dots, m$ contained in a finite connected non-Dynkin quiver Q without oriented cycles and with vertices $1, \dots, n$ ordered as before. Let Λ be the preprojective algebra of Q and $\Lambda_{n+m} = \Lambda / (I_1 \cdots I_n I_1 \cdots I_m)$. Then $\Lambda_m = \Lambda / I_1 \cdots I_m$ is a cluster tilting object in $\underline{\text{Sub}} \Lambda_{n+m}$ with $\text{End}_{\underline{\text{Sub}} \Lambda_{n+m}}(\Lambda_m) \simeq kQ'$.*

Proof. Since we have seen in Lemma II.3.1 that the product $(I_1 \cdots I_n)^2$ gives rise to a strict descending chain of ideals, it follows that the same holds for $I_1 \cdots I_n I_1 \cdots I_m$. The assertions follow as in the proof of Proposition II.3.2. $\square$

Recall from [KR2] that if a connected algebraic triangulated 2-CY category has a cluster tilting object M whose quiver Q has no oriented cycles, then $\mathcal{C}$ is triangle equivalent to the cluster category $\mathcal{C}_{kQ}$. Then we get the following consequence of the last two results.

Theorem II.3.4. *Let Q' be a finite connected quiver without oriented cycles. Let $Q = Q'$ if Q' is not Dynkin, and Q as in Proposition II.3.3 if Q' is Dynkin. Let Λ be the preprojective*

algebra of Q . Then there is a tilting ideal I in Λ such that $\underline{\text{Sub}}\Lambda/I$ is triangle equivalent to the cluster category $\mathcal{C}_{kQ'}$ of Q' .

We finally show that also the categories $\underline{\text{mod}}\Lambda'$ where Λ' is the preprojective algebra of a Dynkin quiver Q' , can be realized this way.

Theorem II.3.5. *Let Q' be a Dynkin quiver contained in a finite connected non-Dynkin quiver Q without loops. We denote by Λ' the preprojective algebra of Q' , by W' the subgroup of W generated by $\{s_i \mid i \in Q'_0\}$, and by w_0 the longest element in W' . Then Λ' is isomorphic to Λ/I_{w_0} and $\underline{\text{mod}}\Lambda' = \underline{\text{Sub}}\Lambda/I_{w_0}$.*

Proof. Let $I_{Q'} := \Lambda(\sum_{i \in Q_0 \setminus Q'_0} e_i)\Lambda$. Since we have $\Lambda/I_{Q'} \simeq \Lambda'$, we only have to show $I_{w_0} = I_{Q'}$. We use the fact that $I_{Q'}$ is maximal amongst all two-sided ideals I of Λ such that any composition factor of Λ/I is S_i for some $i \in Q'_0$.

Since w_0 is a product of s_i ($i \in Q'_0$), any composition factor of Λ/I_{w_0} is S_i for some $i \in Q'_0$. Thus we have $I_{w_0} \supseteq I_{Q'}$. On the other hand, since w_0 is the longest element of W' , we have $l(s_i w_0) < l(w_0)$ for any $i \in Q'_0$. By Proposition II.1.10, we have $I_i I_{w_0} = I_{w_0}$ for any $i \in Q'_0$. This implies $I_{w_0} = I_{Q'}$. $\square$

Using Theorem II.3.5, we see that our theory also applies to preprojective algebras of Dynkin type. In particular, we can specialize Theorem II.2.8 to recover the following result from [GLS1].

Corollary II.3.6. *For a preprojective algebra Λ' of a Dynkin quiver the number of non-isomorphic indecomposable summands in a cluster tilting object is equal to the length $l(w_0)$ of the longest element in the associated Weyl group, which is equal to the number of positive roots.*

We also obtain a large class of cluster tilting objects associated with the different reduced expressions of w_0 .

Our results can also be viewed as giving an interpretation in terms of tilting theory of some functors $\mathcal{E}_i$ used in [GLS5, 5.1].

II.4. Quivers of cluster tilting subcategories.

In this section we show that the quivers of standard cluster tilting subcategories associated with a reduced expression can be described directly from the reduced expression.

Let $s_{i_1} s_{i_2} \cdots s_{i_k} \cdots$ be a (finite or infinite) reduced expression associated with a graph Δ with vertices $1, \dots, n$. We associate with this sequence a quiver $Q(i_1, i_2, \dots)$ as follows, where the vertices correspond to the s_{i_k} .

- For two consecutive i ($i \in \{1, \dots, n\}$), draw an arrow from the second one to the first one.
- For each edge $i - j$, pick out the expression consisting of the i_k which are i or j , so that we have $\cdots ii \cdots jj \cdots jii \cdots i \cdots$. We draw d_{ij} arrows from the last i in a connected set of i 's to the last j in the next set of j 's, and the same from j to i . (Note that since by assumption both i and j occur an infinite number of times if the expression is infinite, each connected set of i 's or j 's is finite.)

Note that in the Dynkin case essentially the same quiver has been used in [BFZ].

For a finite reduced expression $s_{i_1} \cdots s_{i_k}$ we denote by $\underline{Q}(i_1, \dots, i_k)$ the quiver obtained from $Q(i_1, \dots, i_k)$ by removing the last i for each i in Q_0 .

We denote by $\Lambda = T_0 \supsetneq T_1 \supsetneq \cdots$ the associated strict descending chain of tilting ideals. Then we have a cluster tilting subcategory $\mathcal{M}(i_1, i_2, \dots) = \text{add}\{\Lambda_k \mid k > 0\}$ for $\Lambda_k := \Lambda/T_k$.


Theorem II.4.1. *Let the notation be as above.*

- (a) *The quiver of the cluster tilting subcategory $\mathcal{M}(i_1, i_2, \dots)$ is $\underline{Q}(i_1, i_2, \dots)$.*


(b) The quiver of $\underline{\text{End}}_\Lambda(\mathcal{M}(i_1, \dots, i_k))$ is $\underline{Q}(i_1, \dots, i_k)$.

Before we give the proof, we give some examples and consequences.


It follows from the definition that we get the same quiver if we interchange two neighbors in the expression of w which are not connected with any edge in Δ . But if we take two reduced expressions in general, we may get different quivers, as the following examples show.

Let Δ be the graph  and $w = s_1 s_2 s_1 s_3 s_2 = s_2 s_1 s_2 s_3 s_2 = s_2 s_1 s_3 s_2 s_3$


expressions which are clearly reduced. The first expression gives the quiver



the second one gives the quiver



and the third one gives the quiver



We now investigate the relationship between the cluster tilting objects given by different reduced expressions of the same element.

Lemma II.4.2. Let $w = s_{i_1} \cdots s_{i_m} = s_{i'_1} \cdots s_{i'_m}$ be reduced expressions and $\Lambda = T_0 \supsetneq T_1 \supsetneq \cdots$ and $\Lambda = T'_0 \supsetneq T'_1 \supsetneq \cdots$ corresponding tilting ideals.

- (a) Assume that for some k we have $i_k = i'_{k+1}$, $i'_k = i_{k+1}$ and $i_j = i'_j$ for any $j \neq k, k+1$. Then the corresponding cluster tilting objects are isomorphic.
- (b) Assume that for some k we have $i_{k-1} = i'_k = i_{k+1}$, $i'_{k-1} = i_k = i'_{k+1}$ and $i_j = i'_j$ for any $j \neq k, k \pm 1$. Then the corresponding cluster tilting objects are in the relationship of exchanges of $T_{k-1}e_{i_{k-1}}$ and $T'_{k-1}e_{i'_{k-1}}$.

Proof. (a) Obviously we have $T_j = T'_j$ for any $j < k$. Since $s_{i_k} s_{i_{k+1}} = s_{i'_k} s_{i'_{k+1}}$, we have $I_{i_k} I_{i_{k+1}} = I_{i'_k} I_{i'_{k+1}}$. Thus we have $T_j = T'_j$ for any $j > k+1$. In particular, we have $T_j e_{i_j} = T'_j e_{i'_j}$ for any $j \neq k, k+1$. Since $I_{i_k} e_{i_k} = I_{i'_k} I_{i'_{k+1}} e_{i'_{k+1}}$, we have $T_k e_{i_k} = T'_{k+1} e_{i'_{k+1}}$. Similarly, we have $T_{k+1} e_{i_{k+1}} = T'_k e_{i'_k}$. Thus the assertion follows.

(b) Since $I_{i_{k-1}} I_{i_k} I_{i_{k+1}} = I_{i'_{k-1}} I_{i'_k} I_{i'_{k+1}}$, we have $T_j e_{i_j} = T'_j e_{i'_j}$ for any $j \neq k, k \pm 1$. Since $I_{i_{k-1}} I_{i_k} e_{i_k} = I_{i'_{k-1}} I_{i'_k} I_{i'_{k+1}} e_{i'_{k+1}}$, we have $T_k e_{i_k} = T'_{k+1} e_{i'_{k+1}}$. Similarly we have $T_{k+1} e_{i_{k+1}} = T'_k e_{i'_k}$. Thus we have the assertion. $\square$

As an illustration, note that in the above example we obtain the second quiver from the first by mutation at the left vertex. Immediately we have the following conclusion.

Proposition II.4.3. All cluster tilting objects in $\text{Sub}(\Lambda/I_w)$ obtained from reduced expressions of w can be obtained from each other under repeated exchanges.

Proof. This is immediate from Lemma II.4.2 and [BB, 3.3.1] since we get from one reduced expression to another by applying the operations described in Lemma II.4.2. $\square$

Using Theorem II.3.5 we see that for preprojective algebras of Dynkin quivers we get the quivers of the endomorphism algebras associated with reduced expressions of the longest element w_0 .

For a stably 2-CY category or a triangulated 2-CY category $\mathcal{C}$ with cluster tilting subcategories we have an associated *cluster tilting graph* defined as follows. The vertices correspond to the non-isomorphic basic cluster tilting objects, and two vertices are connected with an edge if the corresponding cluster tilting objects differ in exactly one indecomposable summand. For cluster categories this graph is known to be connected [BMRRT], while this is an open problem in general. For the categories $\text{Sub } \Lambda / I_w$ or $\underline{\text{Sub}} \Lambda / I_w$ it follows from Proposition II.4.3 that all standard cluster tilting objects belong to the same component of the cluster tilting graph, and we call this the *standard component*.

We now illustrate with some classes of examples. Let Q be a connected non-Dynkin quiver without oriented cycles, and vertices $1, \dots, n$, where there is no arrow $i \rightarrow j$ for $j > i$.

(a) Let $w = s_1 s_2 \cdots s_n s_1 s_2 \cdots s_n$. The last n vertices correspond to projectives, so the quiver for the cluster tilting object in the stable category $\underline{\text{Sub}} \Lambda / I_w$ is Q , which has no oriented cycles. So we get an alternative proof of Proposition II.3.2.

(b) Choose $w = s_1 s_2 \cdots s_n s_1 s_2 \cdots s_n \cdots$. Ordering the indecomposable preprojective modules as $P_1, \dots, P_n, \tau^{-1} P_1, \dots, \tau^{-1} P_n, \dots, \tau^{-i} P_1, \dots, \tau^{-i} P_n, \dots$ where P_i is the projective module associated with vertex i , we have a bijection between the indecomposable preprojective modules and the terms in the expression for w . Then the quiver of the corresponding cluster tilting subcategory is the preprojective component of the AR quiver of kQ , with an additional arrow from X to τX for each indecomposable preprojective module X . This is a direct consequence of our rule, since we know by Lemma II.3.1 that the expression for w is reduced.


(c) Now take a part $\mathcal{P}$ of the AR quiver of the preprojective component, closed under predecessors. Consider the expression obtained from $s_1 s_2 \cdots s_n s_1 s_2 \cdots s_n \cdots$ by only keeping the terms corresponding to the objects in $\mathcal{P}$ under our given bijection. We show below that this new expression is reduced. Then it follows directly by our rule that when adding arrows $X \rightarrow \tau X$ when X is nonprojective in $\mathcal{P}$, we get the quiver of the cluster tilting object given by the above reduced expression. That this quiver is the quiver of a cluster tilting object was also shown in [GLS3] for $\mathcal{P}$ being the AR quiver of a Dynkin quiver, and in [GLS6] in the general case.

Lemma II.4.4. *The word associated with $\mathcal{P}$ obtained in this way is reduced.*

Proof. The word satisfies the following conditions.

- (a) For each pair (i, j) of vertices connected by some edge, i and j occur each second time after removing the other vertices.
- (b) $w = A_1 A_2 \cdots A_t$, where each A_s is a strict increasing sequence of numbers in $\{1, \dots, n\}$, such that, if $j \notin A_s$, then $j \notin A_{s+1}$, and if $i < j$ are connected with an edge and $i \notin A_s$ (respectively, $j \notin A_s$), then $j \notin A_s$ (respectively, $i \notin A_{s+1}$).

The condition (a) is immediate from the construction. For each pair (i, j) connected with some edge, we have in the AR quiver



the part involving i 's and j 's. Hence they must occur each second time to give this quiver. Thus the condition (a) is satisfied.

If $A_1 \neq (1, \dots, n)$, then w is a subsequence of $(1, \dots, n)$. So the word is clearly reduced in this case. Thus we can assume $A_1 = (1, \dots, n)$. We show that, for any word satisfying the conditions (a)(b) and $A_1 = (1, \dots, n)$, the corresponding descending chain $\Lambda = T_0 \supseteq T_1 \supseteq \cdots$ is strict. Then the word is reduced by Proposition II.1.11.

We assume $T_{k-1} = T_k$ for some k . So $I_{i_1} \cdots I_{i_k} = I_{i_1} \cdots I_{i_{k-1}}$. Take s minimal such that $A_s \neq (1, \dots, n)$, and take i minimal such that $i \notin A_s$. By assumption (b), all terms appearing after the position of i in A_s are not connected with i by an edge in Δ . In

particular, the corresponding ideals commute with I_i . By multiplying with I_i from the right and using commutative relations, we get an equality where i is inserted in A_s after $1, \dots, i-1$. Repeating this process, we get an equality

$$I_1 \cdots I_n I_1 \cdots I_n \cdots I_1 \cdots I_{i_k} = I_1 \cdots I_n I_1 \cdots I_n \cdots I_1 \cdots I_{i_k-1}.$$

This contradicts Lemma II.3.1. $\square$

In the rest of this section we give a proof of Theorem II.4.1. Note that (b) follows directly from (a) and Proposition II.1.11(b).

(a) Let J be the Jacobson radical of Λ . Let $\mathcal{M} = \mathcal{M}(i_1, i_2, \dots)$ and $T_{l,k} = I_{i_l} I_{i_{l+1}} \cdots I_{i_k}$. For $l > k$, this means $T_{l,k} = \Lambda$. In the rest we often use the following equalities.

$$e_i J e_{i'} = \begin{cases} e_i I_i e_{i'} & (i = i') \\ e_i \Lambda e_{i'} & (i \neq i'), \end{cases} \quad I_i e_{i'} = \begin{cases} J e_{i'} & (i = i') \\ \Lambda e_{i'} & (i \neq i'), \end{cases} \quad \text{and} \quad e_{i'} I_i = \begin{cases} e_{i'} J & (i = i') \\ e_{i'} \Lambda & (i \neq i'). \end{cases}$$

We have

$$\text{Hom}_\Lambda(\Lambda_l e_{i_l}, \Lambda_k e_{i_k}) = e_{i_l} (T_{l+1,k}/T_k) e_{i_k}$$

by Lemma II.1.14. We have

$$\text{rad}_\mathcal{M}(\Lambda_l e_{i_l}, \Lambda_k e_{i_k}) = e_{i_l} (T_{(l+1-\delta_{l,k}),k}/T_k) e_{i_k}.$$

Moreover, we have

$$\text{rad}_\mathcal{M}^2(\Lambda_l e_{i_l}, \Lambda_k e_{i_k}) = e_{i_l} ((T_k + \sum_{j>0} T_{(l+1-\delta_{l,j}),j} e_{i_j} T_{(j+1-\delta_{j,k}),k})/T_k) e_{i_k}.$$

To get the quiver of $\mathcal{M}$, we have to compute $(\text{rad}_\mathcal{M} / \text{rad}_\mathcal{M}^2)(\Lambda_l e_{i_l}, \Lambda_k e_{i_k}) = E_{l,k}/D_{l,k}$ for

$$E_{l,k} = e_{i_l} T_{(l+1-\delta_{l,k}),k} e_{i_k} \supseteq D_{l,k} = e_{i_l} T_k e_{i_k} + \sum_{j>0} e_{i_l} T_{(l+1-\delta_{l,j}),j} e_{i_j} T_{(j+1-\delta_{j,k}),k} e_{i_k}.$$

We denote by k^+ the minimal number satisfying $k < k^+$ and $i_k = i_{k^+}$ if it exists.

(i) We consider the case when there are no arrows in Q from l to k . We shall show $E_{l,k} = D_{l,k}$.

If $l > k$ and $i_l = i_k$, then we have $l > k^+ > k$. Thus

$$E_{l,k} = e_{i_l} \Lambda e_{i_k} = e_{i_l} \Lambda e_{i_{k^+}} \Lambda e_{i_k} \subseteq D_{l,k}.$$

In the rest we assume that either $l \leq k$ or $i_l \neq i_k$ holds. First we show

$$E_{l,k} = e_{i_l} T_{l+1,k-1} (1 - e_{i_k}) \Lambda e_{i_k} = \sum_{a \neq i_k} e_{i_l} T_{l+1,k-1} e_a \Lambda e_{i_k}$$

by the following case by case study.

- If $l < k$, then $E_{l,k} = e_{i_l} T_{l+1,k-1} I_{i_k} e_{i_k} = e_{i_l} T_{l+1,k-1} (1 - e_{i_k}) \Lambda e_{i_k}$.
- If $l = k$, then $E_{l,k} = e_{i_l} I_{i_k} e_{i_k} = e_{i_l} \Lambda (1 - e_{i_k}) \Lambda e_{i_k}$.
- If $l > k$ and $i_l \neq i_k$, then $E_{l,k} = e_{i_l} \Lambda e_{i_k} = e_{i_l} I_{i_k} e_{i_k} = e_{i_l} \Lambda (1 - e_{i_k}) \Lambda e_{i_k}$.

Thus we only have to show that $e_{i_l} T_{l+1,k-1} e_a \Lambda e_{i_k} \subseteq D_{l,k}$ for any $a \neq i_k$. We have the following three possibilities.

- If $a \notin \{i_1, i_2, \dots, i_{k-1}\}$, then we have $T_{l+1,k-1} e_a = I_{i_{l+1}} \cdots I_{i_{k-1}} e_a = \Lambda e_a = I_{i_1} \cdots I_{i_{k-1}} e_a = T_{k-1} e_a$. Thus

$$e_{i_l} T_{l+1,k-1} e_a \Lambda e_{i_k} = e_{i_l} T_{k-1} e_a \Lambda e_{i_k} \subseteq e_{i_l} T_k e_{i_k} \subseteq D_{l,k}.$$

- If $a \notin \{i_{k+1}, i_{k+2}, \dots, i_{k^+}\}$, then we have $e_a \Lambda = e_a I_{i_k} \cdots I_{i_{k^+}} = e_a T_{k,k^+}$. Thus

$$e_{i_l} T_{l+1,k-1} e_a \Lambda e_{i_k} = e_{i_l} T_{l+1,k-1} e_a T_{k,k^+} e_{i_k} \subseteq e_{i_l} T_{l+1,k^+} e_{i_k} = e_{i_l} T_{l+1,k^+} e_{i_{k^+}} \Lambda e_{i_k} \subseteq D_{l,k}$$

since $l \neq k^+ \neq k$.

- Otherwise, there is an arrow $j \rightarrow k$ with $i_j = a$. Since $a \notin \{i_{j+1}, i_{j+2}, \dots, i_{k-1}\}$, we have $T_{j+1,k-1} e_a \Lambda = I_{i_{j+1}} \cdots I_{i_{k-1}} e_a \Lambda = \Lambda e_a \Lambda = \Lambda e_a I_{i_{j+1}} \cdots I_{i_{k-1}} = \Lambda e_a T_{j+1,k-1}$. Thus

$$e_{i_l} T_{l+1,k-1} e_a \Lambda e_{i_k} = e_{i_l} T_{l+1,j} T_{j+1,k-1} e_a \Lambda e_{i_k} = e_{i_l} T_{l+1,j} e_{i_j} T_{j+1,k} e_{i_k} \subseteq D_{l,k}$$

since $l \neq j \neq k$.

In each case we have $e_{i_l} T_{l+1,k-1} e_a \Lambda e_{i_k} \subseteq D_{l,k}$ for any $a \neq i_k$.

- (ii) We consider the case $l = k^+$. We have $E_{k^+,k} = e_{i_{k^+}} \Lambda e_{i_k}$. We shall show $D_{k^+,k} = e_{i_l} J e_{i_k}$.

Clearly we have $D_{k^+,k} \subseteq e_{i_{k^+}} J e_{i_k}$. Conversely, we have

$$e_{i_{k^+}} J e_{i_k} = e_{i_{k^+}} I_{i_k} e_{i_k} = e_{i_{k^+}} \Lambda (1 - e_{i_k}) \Lambda e_{i_k} = \sum_{a \neq i_k} e_{i_{k^+}} \Lambda e_a \Lambda e_{i_k}.$$

Thus we only have to show $e_{i_{k^+}} \Lambda e_a \Lambda e_{i_k} \subseteq D_{k^+,k}$ for any $a \neq i_k$. We have the following two possibilities.

- If $a \notin \{i_1, i_2, \dots, i_{k-1}\}$, then we have $\Lambda e_a = I_{i_1} \cdots I_{i_{k-1}} e_a = T_{k-1} e_a$. Thus

$$e_{i_{k^+}} \Lambda e_a \Lambda e_{i_k} = e_{i_{k^+}} T_{k-1} e_a I_{i_k} e_{i_k} \subseteq e_{i_{k^+}} T_k e_{i_k} \subseteq D_{k^+,k}.$$

- If $a \in \{i_1, i_2, \dots, i_{k-1}\}$, then take the largest j such that $i_j = a$. Then we have $\Lambda = T_{k^++1,j}$ and $e_a \Lambda = e_a I_{i_{j+1}} \cdots I_{i_k} = e_a T_{j+1,k}$. Thus

$$e_{i_{k^+}} \Lambda e_a \Lambda e_{i_k} = e_{i_{k^+}} T_{k^++1,j} e_{i_j} T_{j+1,k} e_{i_k} \subseteq D_{k^+,k}$$

since $k^+ \neq j \neq k$.

In each case we have $e_{i_{k^+}} \Lambda e_a \Lambda e_{i_k} \subseteq D_{k^+,k}$ for any $a \neq i_k$.

- (iii) Finally we consider the case when $l \neq k^+$ and there is an arrow in Q from l to k . Then $l < k$. We have $E_{l,k} = e_{i_l} J e_{i_k}$. We shall show $D_{l,k} = e_{i_l} J^2 e_{i_k}$.

First we show $D_{l,k} \subseteq e_{i_l} J^2 e_{i_k}$. We have $e_{i_l} T_k e_{i_k} \subseteq e_{i_l} I_{i_l} I_{i_k} e_{i_k} = e_{i_l} J^2 e_{i_k}$. We have the following three possibilities.

- Assume $l \leq j \leq k$. Then $e_{i_l} T_{(l+1-\delta_{l,j}),j} e_{i_j} T_{(j+1-\delta_{j,k}),k} e_{i_k} \subseteq e_{i_l} J e_{i_j} J e_{i_k} \subseteq e_{i_l} J^2 e_{i_k}$.
- Assume $k < j$. If $i_j \neq i_k$, then $e_{i_l} T_{l+1,j} e_{i_j} T_{j+1,k} e_{i_k} \subseteq e_{i_l} I_{i_j} e_{i_j} J e_{i_k} \subseteq e_{i_l} J^2 e_{i_k}$. If $i_j = i_k$, then $j \geq k^+$. Since there is an arrow $l \rightarrow k$, we have $i_l \in \{i_{k+1}, i_{k+2}, \dots, i_{k^+-1}\} \subseteq \{i_{l+1}, i_{l+2}, \dots, i_{j-1}\}$. Thus $e_{i_l} T_{l+1,j} e_{i_j} T_{j+1,k} e_{i_k} \subseteq e_{i_l} I_{i_l} I_{i_j} e_{i_j} \Lambda e_{i_k} \subseteq e_{i_l} J^2 e_{i_k}$.
- Assume $l > j$. If $i_j \neq i_l$, then $e_{i_l} T_{l+1,j} e_{i_j} T_{j+1,k} e_{i_k} \subseteq e_{i_l} J e_{i_j} I_k e_{i_k} \subseteq e_{i_l} J^2 e_{i_k}$. If $i_j = i_l$, then $e_{i_l} T_{l+1,j} e_{i_j} T_{j+1,k} e_{i_k} \subseteq e_{i_l} \Lambda e_{i_j} I_{i_l} I_{i_k} e_{i_k} \subseteq e_{i_l} J^2 e_{i_k}$.

Next we show $e_{i_l} J^2 e_{i_k} \subseteq D_{l,k}$. We have

$$e_{i_l} J^2 e_{i_k} = e_{i_l} J (1 - e_{i_k}) \Lambda e_{i_k} = \sum_{a \neq i_k} e_{i_l} J e_a \Lambda e_{i_k}.$$

Thus we only have to show $e_{i_l} J e_a \Lambda e_{i_k} \subseteq D_{l,k}$ for any $a \neq i_k$. We have the following two possibilities.

- If $a \notin \{i_1, i_2, \dots, i_{k-1}\}$, then we have $e_{i_l} J e_a = e_{i_l} \Lambda e_a = e_{i_l} I_{i_1} \cdots I_{i_{k-1}} e_a = e_{i_l} T_{k-1} e_a$. Thus

$$e_{i_l} J e_a \Lambda e_{i_k} = e_{i_l} T_{k-1} e_a I_{i_k} e_{i_k} \subseteq e_{i_l} T_k e_{i_k} \subseteq D_{l,k}.$$

- If $a \in \{i_1, i_2, \dots, i_{k-1}\}$, then take the largest j such that $i_j = a$. Then we have $e_a \Lambda = e_a I_{i_{j+1}} \cdots I_{i_k} = e_a T_{j+1,k}$. Moreover if $j = l$, then $e_{i_l} J e_a = e_{i_l} T_{(l+1-\delta_{l,j}),j} e_a$. If $j \neq l$, then $e_{i_l} J e_a \subseteq e_{i_l} I_{i_{l+1}} \cdots I_{i_j} e_a = e_{i_l} T_{l+1,j} e_a$. Thus

$$e_{i_l} J e_a \Lambda e_{i_k} = e_{i_l} T_{(l+1-\delta_{l,j}),j} e_{i_j} T_{j+1,k} e_{i_k} \subseteq D_{l,k}$$

since $j \neq k$.

In each case we have $e_{i_l} J e_a \Lambda e_{i_k} \subseteq D_{l,k}$ for any $a \neq i_k$. $\square$

II.5. Substructure.

In this section we point out that the work in this chapter gives several illustrations of substructures of cluster structures. We also give some concrete examples of 2-CY categories and their cluster tilting objects, to be applied in Chapter III.

Let $s_{i_1}s_{i_2}\cdots s_{i_t}\cdots$ be an infinite reduced expression which contains each $i \in \{1, \dots, n\}$ an infinite number of times. Let $T_t = I_{i_1}\cdots I_{i_t}$, and $\Lambda_t = \Lambda/T_t$. Recall that for $t < m$, we have $\text{Sub } \Lambda_t \subseteq \text{Sub } \Lambda_m \subseteq \text{f.l. } \Lambda$. We then have the following.

Theorem II.5.1. *Let the notation be as above.*

- (a) $\text{Sub } \Lambda_m$, $\underline{\text{Sub}} \Lambda_m$ and $\text{f.l. } \Lambda$ have a cluster structure using the cluster tilting subcategories with the indecomposable projectives as coefficients.
- (b) For $t < m$, the cluster tilting object $\Lambda_1 \oplus \cdots \oplus \Lambda_t$ in $\text{Sub } \Lambda_t$ can be extended to a cluster tilting object $\Lambda_1 \oplus \cdots \oplus \Lambda_t \oplus \cdots \oplus \Lambda_m$ in $\text{Sub } \Lambda_m$, and determines a substructure of $\text{Sub } \Lambda_m$.
- (c) The cluster tilting object $\Lambda_1 \oplus \cdots \oplus \Lambda_t$ in $\text{Sub } \Lambda_t$ can be extended to the cluster tilting subcategory $\{\Lambda_i \mid i \geq 0\}$ in $\text{f.l. } \Lambda$, and determines a substructure of $\text{f.l. } \Lambda$.

Proof. (a) Since $\text{Sub } \Lambda_m$ and $\underline{\text{Sub}} \Lambda_m$ are stably 2-CY and 2-CY respectively, they have a weak cluster structure. It follows from Proposition I.1.11 that we have no loops or 2-cycles, using the cluster tilting objects. Then it follows from Theorem I.1.6 that we have a cluster structure for $\text{Sub } \Lambda_m$ and $\underline{\text{Sub}} \Lambda_m$.

That also $\text{f.l. } \Lambda$ has a cluster structure follows by using that it is the case for all the $\text{Sub } \Lambda_m$.

(b) and (c) follow directly from the definition of substructure and previous results. $\square$

We now consider the Kronecker quiver $1 \rightrightarrows 0$, and let Λ be the associated preprojective algebra. The only strict descending chains are

$$I_0 \supsetneq I_0 I_1 \supsetneq I_0 I_1 I_0 \supsetneq \cdots (I_0 I_1)^j \supsetneq (I_0 I_1)^j I_0 \supsetneq \cdots \quad \text{and} \\ I_1 \supsetneq I_1 I_0 \supsetneq I_1 I_0 I_1 \supsetneq \cdots (I_1 I_0)^j \supsetneq (I_1 I_0)^j I_1 \supsetneq \cdots$$


We let T_t be the product of the first t ideals, and $\Lambda_t = \Lambda/T_t$. Both I_0 and I_1 occur an infinite number of times in each chain. The indecomposable projective Λ -modules P_0 and P_1 have the following structure

$$P_0 = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ \vdots & \vdots & \vdots & \vdots \end{pmatrix} \quad P_1 = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ \vdots & \vdots & \vdots & \vdots \end{pmatrix}$$

where radical layer number $2i$ has $2i$ copies of 1 for P_0 and $2i$ copies of 0 for P_1 , and radical layer number $2i+1$ has $2i+1$ copies of 0 for P_0 and $2i+1$ copies of 1 for P_1 . We write $P_{0,t} = P_0/J^t P_0$ and $P_{1,t} = P_1/J^t P_1$. Then it is easy to see that for the chain $I_0 \supsetneq I_0 I_1 \supsetneq \cdots$ we have $\Lambda_1 = \Lambda/I_0 = P_{0,1} = (0)$, $\Lambda_2 = \Lambda/I_0 I_1 = P_{0,1} \oplus P_{1,2} = (0) \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\Lambda_3 = \Lambda/I_0 I_1 I_0 = P_{0,3} \oplus P_{1,2}, \dots$, $\Lambda_{2t} = P_{0,2t+1} \oplus P_{1,2t}$, $\Lambda_{2t+1} = P_{0,2t-1} \oplus P_{1,2t}, \dots$

Note that this calculation also shows that both our infinite chains are strict descending.

It follows from Section 4 (and is also easily seen directly) that the quiver of the cluster tilting subcategory $\{\Lambda_i \mid i \geq 1\}$ is the following:



In particular, we have the cluster tilting object $P_{0,1} \oplus P_{1,2} \oplus P_{0,3}$ for $\text{Sub } \Lambda_3$, where the last two summands are projective. Hence $P_{0,1}$ is a cluster tilting object in $\underline{\text{Sub}} \Lambda_3$. The quiver of the endomorphism algebra consists of one vertex and no arrows. Hence $\underline{\text{Sub}} \Lambda_3$ is equivalent to the cluster category $\mathcal{C}_k$, which has exactly two indecomposable objects. The other one is $JP_{0,3}$, obtained from the exchange sequence $0 \rightarrow JP_{0,3} \rightarrow P_{0,3} \rightarrow P_{0,1} \rightarrow 0$. Note that it is also easy to see directly that there are no other indecomposable rigid nonprojective objects in $\text{Sub } \Lambda_3$.

For Λ_4 , we have the cluster tilting object $P_{0,1} \oplus P_{1,2} \oplus P_{0,3} \oplus P_{1,4}$ for $\text{Sub } \Lambda_4$. Again the last two Λ_4 -modules are projective, so $P_{0,1} \oplus P_{1,2}$ is a cluster tilting object in $\underline{\text{Sub}} \Lambda_4$. The quiver of the endomorphism algebra is $\cdot \rightrightarrows \cdot$, which has no oriented cycles, and hence $\underline{\text{Sub}} \Lambda_4$ is triangle equivalent to the cluster category $\mathcal{C}_{k(\cdot \rightrightarrows \cdot)}$. In particular the cluster tilting graph is connected. We can use this to get a description of the rigid objects in $\text{Sub } \Lambda_4$.

Proposition II.5.2. *Let $\Lambda_4 = \Lambda/I_0I_1I_0I_1$ be the algebra defined above. Then the indecomposable rigid Λ_4 -modules in $\text{Sub } \Lambda_4$ are exactly the ones of the form $\Omega_{\Lambda_4}^i(P_{0,1})$ and $\Omega_{\Lambda_4}^i(P_{1,2})$ for $i \in \mathbb{Z}$*

Proof. For $\mathcal{C}_{k(\cdot \rightrightarrows \cdot)}$ the indecomposable rigid objects are the τ -orbits of the objects induced by the indecomposable projective $k(\cdot \rightrightarrows \cdot)$ -modules. Here $\tau = [1]$, and for $\underline{\text{Sub}} \Lambda_4$, $\Omega^{-1} = [1]$. This proves the claim. $\square$

The cluster tilting graph for $\text{Sub } \Lambda_3$ and $\text{Sub } \Lambda_4$ are $\cdot \text{---} \cdot$ and $\cdots \text{---} \cdot \text{---} \cdot \text{---} \cdot \text{---} \cdot \text{---} \cdots$.

We end with the following problem.

Conjecture II.5.3. *For any $w \in W$ the cluster tilting graph for $\text{Sub } \Lambda/I_w$ is connected.*

III. CONNECTIONS TO CLUSTER ALGEBRAS

While the theory of 2-CY categories is interesting in itself, one of the motivations for investigating 2-CY categories comes from the theory of cluster algebras initiated by Fomin and Zelevinsky [FZ2]. In many situations the 2-CY categories can be used to construct new examples of cluster algebras, and also to give a new categorical model for already known examples. This has been done in for example [CK1, CK2] and [GLS1]. In this chapter we illustrate with some applications of the theory developed in the first two chapters. In Section 2, we recall the definition and basic properties of a map φ from the finite dimensional modules over the completed preprojective algebra of a connected quiver with no loops to the function field $\mathbb{C}(U)$ of the associated unipotent group U . This was used in [GLS1] to model the cluster algebra structure of the coordinate ring $\mathbb{C}[U]$ in the Dynkin case. It is what we call a (strong) cluster map. We also make explicit the notion of subcluster algebra, and observe that a substructure of a (stably) 2-CY category together with a cluster map gives rise to a subcluster algebra. This gives one way to construct new cluster algebras inside $\mathbb{C}[U]$ in the Dynkin case, or model old ones, as we illustrate with examples in Section 2. In Section 3 we deal with the non-Dynkin case. We conjecture that the stably 2-CY category $\text{Sub } \Lambda/I_w$ discussed in Chapter II gives a model for a cluster algebra structure on the coordinate ring $\mathbb{C}[U^w]$ of the corresponding unipotent cell. We give examples which support this conjecture.

III.1. Cluster algebras, subcluster algebras and cluster maps.

In this section we recall the notion of cluster algebras [FZ2] and make explicit a notion of *subcluster algebras*. Actually we extend the definition of cluster algebras to include the possibility of clusters with countably many elements. The coordinate rings of unipotent groups of non-Dynkin diagrams are candidates for containing such cluster algebras. We also

introduce certain maps, called (strong) *cluster maps*, defined for categories with a cluster structure. The image of a cluster map gives rise to a cluster algebra. Cluster substructures on the category side give rise to subcluster algebras.

We first recall the definition of a cluster algebra, allowing countable clusters. Note that the setting used here is not the most general one. Let $m \geq n$ be positive integers, or countable numbers. Let $\mathbf{F} = \mathbb{Q}(u_1, \dots, u_m)$ be the field of rational functions over $\mathbb{Q}$ in m independent variables. A cluster algebra is a subring of $\mathbf{F}$, constructed in the following way. A *seed* in $\mathbf{F}$ is a triple $(\underline{x}, \underline{c}, \tilde{B})$, where $\underline{x}$ and $\underline{c}$ are non-overlapping sets of elements in $\mathbf{F}$, where we let $\tilde{x} = \underline{x} \cup \underline{c}$ and sometimes denote the seed by the pair $(\tilde{x}, \tilde{B})$. Here $\tilde{x} = \{x_1, \dots, x_m\}$ should be a transcendence basis for $\mathbf{F}$ and $\tilde{B} = (b_{ij})$ is a locally finite $m \times n$ -matrix with integer elements such that the submatrix B of $\tilde{B}$ consisting of the n first rows is skew-symmetric.

The set $\underline{x} = \{x_1, \dots, x_n\}$ is called the *cluster* of the seed, and the set $\underline{c} = \{x_{n+1}, \dots, x_m\}$ is the *coefficient set* of the cluster algebra. The set $\tilde{x} = \underline{x} \cup \underline{c}$ is called an *extended cluster*.

For a seed $(\tilde{x}, \tilde{B})$, with $\tilde{B} = (b_{ij})$, and for $k \in \{1, \dots, n\}$, a *seed mutation* in direction k produces a new seed $(\tilde{x}', \tilde{B}')$. Here $\tilde{x}' = (\tilde{x} \setminus \{x_k\}) \cup \{x'_k\}$, where

$$x'_k = x_k^{-1} \left(\prod_{b_{ik} > 0} x_i^{b_{ik}} + \prod_{b_{ik} < 0} x_i^{-b_{ik}} \right)$$

This is called an *exchange relation* and $\{x_k, x'_k\}$ is called an *exchange pair*. Furthermore

$$b'_{ij} = \begin{cases} -b_{ij} & \text{if } i = k \text{ or } j = k \\ b_{ij} + \frac{|b_{ik}|b_{kj} + b_{ik}|b_{kj}|}{2} & \text{else.} \end{cases}$$

Fix an (initial) seed $(\tilde{x}, \tilde{B})$, and consider the set $\mathbf{S}$ of all seeds obtained from $(\tilde{x}, \tilde{B})$ by a sequence of seed mutations. The union $\mathbf{X}$ of all elements in the clusters in $\mathbf{S}$ are called the *cluster variables*, and for a fixed subset of coefficients $\underline{c}_0 \subseteq \underline{c}$, the *cluster algebra* $\mathbf{A}(\mathbf{S})$ with the coefficients $\underline{c}_0$ inverted, is the $\mathbb{Z}[\underline{c}, \underline{c}_0^{-1}]$ -subalgebra of $\mathbf{F}$ generated by $\mathbf{X}$. Note that we, unlike in the original definition, do not necessarily invert all coefficients. This is done in order to catch examples like the coordinate ring of a maximal unipotent group in the Dynkin case and the homogeneous coordinate ring of a Grassmannian. Note that we often extend the scalars for cluster algebras to $\mathbb{C}$.

We now make explicit the notion of subcluster algebras. Let $\mathbf{A}$ be a cluster algebra with cluster variables $\mathbf{X}$, coefficients $\underline{c}$, and ambient field $\mathbf{F} = \mathbb{Q}(u_1, \dots, u_m)$. A *subcluster algebra* $\mathbf{A}'$ of $\mathbf{A}$ is a cluster algebra such that there exists a seed $(\underline{x}, \underline{c}, Q)$ for $\mathbf{A}$ and a seed $(\underline{x}', \underline{c}', Q')$ for $\mathbf{A}'$ such that

- (S1) $\underline{x}' \subseteq \underline{x}$ and $\underline{c}' \subseteq \underline{x} \cup \underline{c}$.
- (S2) For each cluster variable $x_i \in \underline{x}'$, the set of arrows entering and leaving i in Q lie in Q' .
- (S3) The invertible coefficients $\underline{c}'_0 \subseteq \underline{c}'$ satisfy $\underline{c}_0 \cap \underline{c}' \subseteq \underline{c}'_0$.

Note that a subcluster algebra is not necessarily a subalgebra since some coefficients may be inverted. Also note that $\mathbf{A}'$ is determined by the seed $(\underline{x}', \underline{c}', Q')$ and the set $\underline{c}'_0$ of invertible coefficients.

The definition implies that clusters in the subcluster algebra can be uniformly extended.

Proposition III.1.1. (a) *Seed mutation in $\mathbf{A}'$ is compatible with seed mutation in $\mathbf{A}$.*
 (b) *There is a set $\underline{v}$ consisting of cluster variables and coefficients in $\mathbf{A}$, such that for any extended cluster $\underline{x}'$ in $\mathbf{A}'$, $\underline{x}' \cup \underline{v}$ is an extended cluster in $\mathbf{A}$.*

Proof. This follows directly from the definition. $\square$

Inspired by [GLS1, GLS4] and [CC, CK1, CK2] we introduce certain maps, which we call (strong) cluster maps, defined for a 2-CY category with a (weak) cluster structure,

and such that the image gives rise to a cluster algebra. We show that such maps preserve substructures, as defined above and in Section I.2.

Recall that a category $\mathcal{C}$ is stably 2-CY if it is either an exact Frobenius category where $\underline{\mathcal{C}}$ is 2-CY or a functorially finite extension closed subcategory $\mathcal{B}$ of a triangulated 2-CY category $\mathcal{C}$.

Let $\mathcal{C}$ be a stably 2-CY category with a cluster structure defined by cluster tilting objects, where projectives are coefficients. We assume that the cluster tilting objects have n cluster variables and c coefficients, where $1 \leq n \leq \infty$ and $0 \leq c \leq \infty$. For a cluster tilting object T , we denote by $B_{\text{End}_{\mathcal{C}}(T)}$ the $m \times n$ -matrix obtained by removing the last $m - n$ columns of the skew-symmetric $m \times m$ matrix corresponding to the quiver of the endomorphism algebra $\text{End}_{\mathcal{C}}(T)$, where the columns are ordered such that those corresponding to projective summands of T come last. We can also think of this as dropping the arrows between vertices in $\text{End}_{\mathcal{C}}(T)$ corresponding to indecomposable projective summands of T from the quiver of $\text{End}_{\mathcal{C}}(T)$.

Let $\mathbf{F} = \mathbb{Q}(u_1, \dots, u_m)$. Given a connected component Δ of the cluster tilting graph of $\mathcal{C}$, a *cluster map* (respectively, *strong cluster map*) for Δ is a map $\varphi: \mathcal{E} = \text{add}\{T \mid T \in \Delta\} \rightarrow \mathbf{F}$ (respectively, $\varphi: \mathcal{C} \rightarrow \mathbf{F}$) where isomorphic objects have the same image, satisfying the following three conditions.

- (M1) For a cluster tilting object T in Δ , $\varphi(T)$ is a transcendence basis for $\mathbf{F}$.
- (M2) (respectively, (M2')) For all indecomposable objects M and N in $\mathcal{E}$ (respectively, $\mathcal{C}$) with $\dim_k \text{Ext}^1(M, N) = 1$, we have $\varphi(M)\varphi(N) = \varphi(V) + \varphi(V')$ where V and V' are the middle of the non-split triangles/short exact sequences $N \rightarrow V \rightarrow M$ and $M \rightarrow V' \rightarrow N$.
- (M3) (respectively, (M3')) $\varphi(A \oplus A') = \varphi(A)\varphi(A')$ for all A, A' in $\mathcal{E}$ (respectively, $\mathcal{C}$).

Note that a pair (M, N) of indecomposable objects in $\mathcal{E}$ is an exchange pair if and only if $\text{Ext}^1(M, N) \simeq k$ (see [BMRR]). Note that a map $\varphi: \mathcal{C} \rightarrow \mathbf{F}$ satisfying (M2') and (M3') is called a *cluster character* in [P]. Important examples of (strong) cluster maps appear in [CK1, CK2, GLS1], and more recently in [GLS6, P].

Theorem III.1.2. *With $\mathcal{C}$ and $\mathcal{E}$ as above, let $\varphi: \mathcal{E} \rightarrow \mathbf{F}$ be a cluster map. Then the following hold.*

- (a) *Let $\mathbf{A}$ be the subalgebra of $\mathbf{F}$ generated by $\varphi(X)$ for $X \in \mathcal{E}$. Then $\mathbf{A}$ is a cluster algebra and $(\varphi(T), B_{\text{End}_{\mathcal{C}}(T)})$ is a seed for $\mathbf{A}$ for any cluster tilting object T in Δ .*
- (b) *Let $\mathcal{B}$ be a subcategory of $\mathcal{C}$ with a substructure, and $\mathcal{E}'$ a subcategory of $\mathcal{B}$ defined by a connected component of the cluster tilting graph of $\mathcal{B}$. Then $\varphi(X)$ for $X \in \mathcal{E}'$ generates a subcluster algebra of $\mathbf{A}$.*

Proof. (a) follows from the fact that $\mathcal{C}$ has a cluster structure. For (b), let T' be a cluster tilting object in the subcategory that extends to a cluster tilting object T for $\mathcal{E}$. Then $\varphi(T')$ gives a transcendence basis for a subfield $\mathbf{F}'$ of $\mathbf{F}$, and T' together with its matrix $B_{\text{End}_{\mathcal{C}}(T')}$ gives a seed for a subcluster algebra. $\square$

For any subset $\underline{\mathcal{C}}_0$ of coefficients of $\mathbf{A}$, we have a cluster algebra $\mathbf{A}[\underline{\mathcal{C}}_0^{-1}]$. We say that the cluster algebra $\mathbf{A}[\underline{\mathcal{C}}_0^{-1}]$ is modelled by the cluster map $\varphi: \mathcal{E} \rightarrow \mathbf{F}$.

III.2. The GLS φ -map with applications to the Dynkin case.

Let Q be a finite connected quiver without loops, and let Λ be the associated completed pre-projective algebra over an algebraically closed field k , and W the associated Coxeter group. For the non-Dynkin case, we have the derived 2-CY category f.l. Λ of finite dimensional left Λ -modules, and for each w in W the stably 2-CY category $\text{Sub } \Lambda/I_w$ as investigated in Chapter II.

On the other hand, associated with the underlying graph of Q , is a Kac-Moody group G [KP1] with a maximal unipotent subgroup U . Let H be the torus. Recall that the Weyl group $\text{Norm}(H)/H$ is isomorphic to the Coxeter group W associated with Q . For an element $w \in W = \text{Norm}(H)/H$ and any lifting $\tilde{w}$ of w in G , define the unipotent cell U^w [BZ] to be the intersection

$$U^w = U \cap B_- \tilde{w} B_-,$$

where B_- is the opposite Borel subgroup corresponding to U . Then U^w is independent of the choice of lifting of w . It is a quasi-affine algebraic variety of dimension $l(w)$ and we have $U = \bigsqcup_{w \in W} U^w$.

Let $U(\mathfrak{n})$ be enveloping algebra of the maximal nilpotent subalgebra of the Kac-Moody Lie algebra $\mathfrak{g}$ associated to Q and let $U(\mathfrak{n})^*$ and $U(\mathfrak{n})_{\text{gr}}^*$ be the dual and graded dual of $U(\mathfrak{n})$ respectively. Note that both $U(\mathfrak{n})^*$ and $U(\mathfrak{n})_{\text{gr}}^*$ become algebras with respect to the $\circ$ -product which is dual to the coproduct on $U(\mathfrak{n})$.

Recall that the matrix coefficient function $f_{v,\zeta}^\tau$ associated to a representation $\tau : U(\mathfrak{n}) \rightarrow \mathfrak{gl}(V)$ and vectors $v \in V$ and $\zeta \in V^*$ is the linear form in $U(\mathfrak{n})^*$ defined by $f_{v,\zeta}^\tau(x) = \zeta(\tau(x) \cdot v)$.

Define the restricted dual $U(\mathfrak{n})_{\text{res}}^*$ of the enveloping algebra $U(\mathfrak{n})$ to be the span of the unit element $1 \in U(\mathfrak{n})^*$ together with all matrix coefficient functions f_{v,v^*}^τ for integrable lowest weight representations $\tau : U(\mathfrak{n}) \rightarrow \mathfrak{gl}(V)$ with $v \in V$ and $\zeta \in V^*$. The restricted dual $U(\mathfrak{n})_{\text{res}}^*$ is a subalgebra of $U(\mathfrak{n})^*$ due to the fact that $f_{v,\zeta}^\tau \circ f_{v',\zeta'}^{\tau'} = f_{v \otimes v', \zeta \otimes \zeta'}^{\tau \otimes \tau'}$. One can check that the graded dual $U(\mathfrak{n})_{\text{gr}}^*$ is a subalgebra of the restricted dual $U(\mathfrak{n})_{\text{res}}^*$.

Let $\mathbb{C}[U]$ denote the ring of (strongly) regular functions on the unipotent group U as defined in [KP2]. Given a representation $\varrho : U \rightarrow \text{GL}(V)$ of the unipotent group U we may differentiate it to obtain a $U(\mathfrak{n})$ -representation $d\varrho : U(\mathfrak{n}) \rightarrow \mathfrak{gl}(V)$ uniquely determined by the formula

$$d\varrho(x) \cdot v = \left. \frac{\partial}{\partial t} \right|_{t=0} \exp(tx) \cdot v$$

for all $x \in \mathfrak{n}$. Accordingly a representation $\varrho : U \rightarrow \text{GL}(V)$ will be called an integrable lowest weight representation of the unipotent group U if the differentiated representation $d\varrho : U(\mathfrak{n}) \rightarrow \mathfrak{gl}(V)$ is integrable and lowest weight.

Theorem 1 of [KP2] implies that $\mathbb{C}[U]$ is spanned by the unit element together with all matrix coefficient functions $F_{v,\zeta}^\varrho$ for integrable lowest weight representations $\varrho : U \rightarrow \text{GL}(V)$ with $v \in V$ and $\zeta \in V^*$. The function $F_{v,\zeta}^\varrho : U \rightarrow \mathbb{C}$ is strongly regular and is defined for $g \in U$ by $F_{v,\zeta}^\varrho(g) := \zeta(\varrho(g) \cdot v)$. One easily checks that, with regard to the usual product of functions, the identity $F_{v,\zeta}^\varrho F_{v',\zeta'}^{\varrho'} = F_{v \otimes v', \zeta \otimes \zeta'}^{\varrho \otimes \varrho'}$ holds.

In view of these facts it follows that the mapping ι given by

$$f_{v,\zeta}^{d\varrho} \mapsto F_{v,\zeta}^\varrho$$

defines an algebra isomorphism between the restricted dual $U(\mathfrak{n})_{\text{res}}^*$ and the coordinate ring $\mathbb{C}[U]$. In particular ι restricts to an embedding of the graded dual $U(\mathfrak{n})_{\text{gr}}^*$ into $\mathbb{C}[U]$.

We now apply the construction of the GLS φ -map [GLS1] to the non-Dynkin case. For $M \in \text{f.l.}\Lambda$, Geiss-Leclerc-Schröer [GLS6] constructed $\delta_M \in U(\mathfrak{n})_{\text{gr}}^*$ by using Lusztig's Lagrangian construction of $U(\mathfrak{n})$ [Lu1, Lu2]. We denote by $\varphi(M) \in \mathbb{C}[U]$ the image of δ_M under the above map $\iota : U(\mathfrak{n})_{\text{gr}}^* \rightarrow \mathbb{C}[U]$. Then $\varphi(M)$ has the following property.

Let $x_i(a) = \exp(ae_i)$ for $a \in \mathbb{C}$, $i \in Q_0$ be elements in U , for a Chevalley generator e_i . For $\mathbf{i} = (i_1, \dots, i_k) \in \mathbb{Z}_{\geq 0}^k$ the symbol $\chi_{\mathbf{i}}(M)$ denotes the Euler characteristic of the variety

$\mathcal{F}_i(M)$ of all flags in $M = M_0 \supset M_1 \supset \cdots \supset M_k = 0$ with M_{l-1}/M_l isomorphic to the simple module S_{i_l} . Then for any $\mathbf{i} = (i_1, \dots, i_k) \in \mathbb{Z}_{\geq 0}^k$ and $a_1, \dots, a_k \in \mathbb{C}$, we have

$$(5) \quad \varphi(M)(x_{i_1}(a_1) \cdots x_{i_k}(a_k)) = \sum_{\mathbf{j} \in \mathbb{Z}_{\geq 0}^k} \chi_{\text{rev } \mathbf{j}}(M) \frac{a_1^{j_1} \cdots a_k^{j_k}}{j_1! \cdots j_k!}$$

where $\text{rev } \mathbf{j}$ is the $j_1 + \cdots + j_k$ tuple which, when read from left to right, starts with j_k occurrences of i_k , followed by j_{k-1} occurrences of i_{k-1} , and ultimately j_1 occurrences of i_1 .

Notice that one can prove that property (5) uniquely determines $\varphi(M)$. The following result (a) was shown in [GLS4, GLS6] and (b) was shown in [GLS1, GLS2].

Theorem III.2.1. (a) $\varphi : \text{f.l. } \Lambda \rightarrow \mathbb{C}[U] \subset \mathbb{C}(U)$ satisfies $(M\mathcal{J})$ and $(M\mathcal{J}')$.
 (b) If Q is Dynkin, then $\mathbb{C}[U]$ (respectively, $\mathbb{C}[U^{w_0}]$ for the longest element w_0) is a cluster algebra modelled by a strong cluster map $\varphi : \text{mod } \Lambda \rightarrow \mathbb{C}(U)$ for the standard component of the cluster tilting graph of $\text{mod } \Lambda$ with no (respectively, all) coefficients inverted.

The image of a substructure $\mathcal{B}$ of $\text{mod } \Lambda$ gives a subcluster algebra of $\mathbb{C}[U]$ for the Dynkin case, and we illustrate this with the examples from I.3. We omit the calculation involved in proving the isomorphisms between the subcluster algebras arising from $\mathcal{B}$ and the coordinate rings of the varieties under consideration. See [BL] for general background on Schubert varieties and (isotropic) Grassmannians. For a subset J of size k in $[1 \dots n]$ the symbol $[J]$ will denote the $k \times k$ matrix minor of an $n \times n$ matrix with row set $[1 \dots k]$ and column set J . In the first two examples G is $\text{SL}_n(\mathbb{C})$ and U is the subgroup of all upper triangular $n \times n$ unipotent matrices.

Example 1 ($\text{Gr}_{2,5}$ -Schubert variety)

Let Λ be of type A_4 , and let $\mathcal{B}$ be the full additive subcategory of $\text{mod } \Lambda$ from Example 1 in I.3. The associated algebraic group is then $\text{SL}_5(\mathbb{C})$. Consider the Grassmannian $\text{Gr}_{2,5}$, and the Schubert variety $X_{3,5}$ associated with the subset $\{3, 5\}$ of $\{1, 2, 3, 4, 5\}$. Let $w_{3,5} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 4 & 1 & 5 & 2 \end{pmatrix}$ be the associated Grassmann permutation in S_5 , and $U^{w_{3,5}}$ the unipotent cell in U associated to $w_{3,5}$. Note that the Schubert variety $X_{3,5}$ is birationally isomorphic to the unipotent cell $U^{w_{3,5}}$ [BZ]. Then $\mathbb{C}[U^{w_{3,5}}]$ is known to be a subcluster algebra of $\mathbb{C}[U]$ [BFZ].

Under the GLS-map φ from $\text{mod } \Lambda$ to $\mathbb{C}[U]$ one can check that $\varphi(M_x) = [x]$, with M_x as defined in Example 1 in I.3. Since $\mathcal{B}$ has a cluster substructure of $\text{mod } \Lambda$, we know that the image gives rise to a subcluster algebra of $\mathbb{C}[U]$. Then the image of $\mathcal{B}$ under the strong cluster map φ is precisely $\mathbb{C}[U^{w_{3,5}}]$. To see this, we mutate a seed from [BFZ] which generates the cluster algebra structure for $\mathbb{C}[U]$, to get a new seed which contains $\varphi(T)$ for the cluster tilting object T in $\mathcal{B}$ in Example 1 in I.3. Then one proves that the image is $\mathbb{C}[U^{w_{3,5}}]$ after a proper choice of which coefficients to invert.

Example 2 (Unipotent Cell in $\text{SL}_4(\mathbb{C})$)

Let Λ be the preprojective algebra of type A_3 , and $\mathcal{B}$ the subcategory of $\text{mod } \Lambda$ from Example 2 in I.3. The associated algebraic group is $\text{SL}_4(\mathbb{C})$. Let U^w be the unipotent cell of the unipotent subgroup U , associated with the permutation $w = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 3 & 1 & 2 \end{pmatrix}$. It is shown in [BFZ] that $\mathbb{C}[U^w]$ is a cluster algebra of type A_2 , and implicitly that it is a subcluster algebra of $\mathbb{C}[U]$. In view of Section III.1, the image $\varphi(\mathcal{B})$ has a subcluster algebra structure modelled by $\mathcal{B}$. As in Example 1, one begins by mutating a seed from [BFZ] which generates the cluster algebra structure for $\mathbb{C}[U]$, such that the new seed contains $\varphi(T)$. Then one proves that the image is $\mathbb{C}[U^w]$, after a proper choice of which coefficients to invert.

Example 3 (The $\mathrm{SO}_8(\mathbb{C})$ -Isotropic Grassmannians (cf. [GLS5, 10.4.3]))

Let Λ be the preprojective algebra of the Dynkin quiver D_4 . Let ϱ be the 4×4 anti-diagonal matrix whose i, j entry is $(-1)^i \delta_{i, 5-j}$ and let J be the 8×8 anti-diagonal matrix, written in block form as

$$\begin{pmatrix} 0 & \varrho \\ \varrho^T & 0 \end{pmatrix}.$$

The *even special orthogonal group* $\mathrm{SO}_8(\mathbb{C})$ is the group of 8×8 matrices

$$\left\{ g \in \mathrm{SL}_8(\mathbb{C}) \mid g^T J g = J \right\}.$$

The *maximal unipotent subgroup* U of $\mathrm{SO}_8(\mathbb{C})$ consists of all 8×8 matrices in $\mathrm{SO}_8(\mathbb{C})$ which are upper triangular and unipotent, i.e. having all diagonal entries equal to 1. A more explicit description in terms of matrices in block form is

$$(6) \quad U = \left\{ \begin{pmatrix} u & u\varrho v \\ 0 & \varrho^T(u^{-1})^T \varrho \end{pmatrix} \mid \begin{array}{l} u \text{ is upper triangular unipotent in } \mathrm{SL}_4(\mathbb{C}) \\ v \text{ is skew-symmetric in } \mathrm{M}_4(\mathbb{C}) \end{array} \right\}$$

The isotropic Grassmannian $\mathrm{Gr}_{2,8}^{\mathrm{iso}}$ is the closed subvariety of the classical Grassmannian $\mathrm{Gr}_{2,8}$ consisting of all isotropic 2-dimensional subspaces of $\mathbb{C}^8$. Let $\widehat{\mathrm{Gr}}_{2,8}^{\mathrm{iso}}$ be the corresponding affine cone. Let $q: U \rightarrow \widehat{\mathrm{Gr}}_{2,8}^{\mathrm{iso}}$ denote the map given by $q(u) = u_1 \wedge u_2$, where u_1, u_2 are the first two rows of u in U , and let $q^*: \mathbb{C}[\widehat{\mathrm{Gr}}_{2,8}^{\mathrm{iso}}] \rightarrow \mathbb{C}[U]$ be the associated homomorphism of coordinate rings.

Let $\varphi: \mathrm{mod} \Lambda \rightarrow \mathbb{C}[U]$ be the GLS φ -map. Then one can show that

$$\begin{array}{llllll} \varphi(M_{16}) &= [16] & \varphi(M_{24}) &= [24] & \varphi(M_{25}) &= [25] & \varphi(M_{26}) &= [26] \\ \varphi(M_{68}) &= [68] & \varphi(M_{18}) &= [18] & \varphi(M_-) &= \psi_- & \varphi(M_+) &= \psi_+ \\ \varphi(P_1) &= [8] & \varphi(P_2) &= [78] & \varphi(P_3) &= \frac{[678]}{\mathrm{Pfaff}_{[1234]}} & \varphi(P_4) &= \mathrm{Pfaff}_{[1234]} \end{array}$$

Here $\mathrm{Pfaff}_{[1234]}$ denotes the Pfaffian of the 4×4 skew-symmetric part v appearing in (6) of the unipotent element and $\psi_{\pm} = \frac{1}{2}([18] - [27] + [36] \pm [45])$. The functions $\frac{[678]}{\mathrm{Pfaff}_{[1234]}}$ and $\mathrm{Pfaff}_{[1234]}$ are examples of generalized minors of type D [FZ1].

In the notation of Example 3 in Section I.3, we have seen that

$$T = M_{16} \oplus M_{24} \oplus M_{25} \oplus M_{26} \oplus M_{68} \oplus M_{18} \oplus M_- \oplus M_+ \oplus P_2$$

is a cluster tilting object in $\mathcal{B} = \mathrm{Sub} P_2$, which can be extended to a cluster tilting object

$$\tilde{T} = T \oplus P_1 \oplus P_3 \oplus P_4$$

for $\mathrm{mod} \Lambda$. One shows that the initial seed used in [BFZ], which determines a cluster algebra structure for $\mathbb{C}[U]$, is mutation equivalent to the initial seed determined by $\tilde{T}$, which hence generates the same cluster algebra. Since the subcategory $\mathcal{B}$ of $\mathrm{mod} \Lambda$ in I.3, Example 3, has a substructure, as defined in Chapter I, the connected component of the cluster tilting graph of $\mathcal{B}$ containing T determines a subcluster algebra $\mathbf{A}'$ of $\mathbb{C}[U]$ (where $[18], \psi_{\pm}, [78]$ are taken as noninverted coefficients). Then we can prove that $\mathbf{A}'$ coincides with $\mathrm{Im} q^*$ (which we conjecture to be true more generally).

Notice that we have the cluster algebra structure for $\widehat{\mathrm{Gr}}_{2,8}^{\mathrm{iso}}$ by adjoining the coefficient $[12]$ to $\mathrm{Im} q^*$.

III.3. Cluster Structure of the loop group $\mathrm{SL}_2(\mathcal{L})$.

Let Q be a finite connected non-Dynkin quiver without loops, and let Λ be the associated completed preprojective algebra and W the associated Coxeter group. Let G be the associated Kac-Moody group G with a maximal unipotent subgroup U , and let U^w be the unipotent cell associated with $w \in W$. Using the GLS-map $\varphi : \mathrm{f.l.}\Lambda \rightarrow \mathbb{C}[U]$ and the restriction map $\mathbb{C}[U] \rightarrow \mathbb{C}[U^w]$, define the induced map

$$\varphi_w : \mathrm{Sub}\Lambda/I_w \subset \mathrm{f.l.}\Lambda \xrightarrow{\varphi} \mathbb{C}[U] \rightarrow \mathbb{C}[U^w].$$

Using our results from Chapter II we know that the transcendence degree $l(w)$ of $\mathbb{C}(U^w)$ is equal to the number of non-isomorphic summands of a cluster tilting object in $\mathrm{Sub}\Lambda/I_w$. It is then natural to pose the following.

Conjecture III.3.1. *For any $w \in W$, the coordinate ring $\mathbb{C}[U^w]$ is a cluster algebra modelled by a strong cluster map $\varphi_w : \mathrm{Sub}\Lambda/I_w \rightarrow \mathbb{C}[U^w]$ for the standard component of the cluster tilting graph of $\mathrm{Sub}\Lambda/I_w$ with all coefficients inverted.*

Recall that any infinite reduced expression where all generators occur an infinite number of times gives rise to a cluster tilting subcategory with an infinite number of non-isomorphic indecomposable objects. Since the GLS-map $\varphi : \mathrm{f.l.}\Lambda \rightarrow \mathbb{C}[U]$ satisfies (M2') and (M3'), it is natural to ask the following.

Question III.3.2. *The coordinate ring $\mathbb{C}[U]$ contains a cluster algebra modelled by $\varphi : \mathrm{f.l.}\Lambda \rightarrow \mathbb{C}[U]$ for any connected component of the cluster tilting graph of $\mathrm{f.l.}\Lambda$.*

As a support for Conjecture III.3.1, we show that this is the case when Q is the Kronecker quiver $1 \rightrightarrows 0$, and the length of w is at most 4. Without loss of generality, we only have to consider the case $w = w_i$ for $w_1 = s_0$, $w_2 = s_0 s_1$, $w_3 = s_0 s_1 s_0$ and $w_4 = s_0 s_1 s_0 s_1$. The cases w_i for $i = 1$ or 2 are clear since Λ/I_{w_i} is a cluster tilting object in $\mathrm{Sub}\Lambda/I_{w_i}$ and $\mathbb{C}[U^{w_i}]$ is generated by invertible coefficients. For the case w_i for $i = 3$ or 4 , we have cluster tilting objects $T_3 = P_{0,1} \oplus P_{1,2} \oplus P_{0,3}$ in $\mathrm{Sub}\Lambda/I_{w_3}$ and $T_4 = P_{0,1} \oplus P_{1,2} \oplus P_{0,3} \oplus P_{1,4}$ in $\mathrm{Sub}\Lambda/I_{w_4}$, where $P_{i,k} = P_i/J^k P_i$ for $i = 0, 1$ and $k > 0$.

The Kac-Moody group $\widehat{\mathrm{SL}}_2(\mathcal{L})$ associated with the Kronecker quiver is defined as the unique non-trivial central extension

$$1 \longrightarrow \mathbb{C}^* \longrightarrow \widehat{\mathrm{SL}}_2(\mathcal{L}) \xrightarrow{\pi} \mathrm{SL}_2(\mathcal{L}) \longrightarrow 1$$

of the algebraic loop group $\mathrm{SL}_2(\mathcal{L})$ by $\mathbb{C}^*$ (see [KP1, PS] for details.) The group $\mathrm{SL}_2(\mathcal{L})$ consists of all $\mathcal{L}$ -valued 2×2 matrices $g = (g_{ij})$ with determinant 1, where $\mathcal{L}$ is the Laurent polynomial ring $\mathbb{C}[t, t^{-1}]$. The maximal unipotent subgroup (respectively, unipotent cells) of $\widehat{\mathrm{SL}}_2(\mathcal{L})$ is mapped isomorphically onto the maximal unipotent subgroup

$$U = \left\{ g \in \begin{pmatrix} 1 + t\mathbb{C}[t] & \mathbb{C}[t] \\ t\mathbb{C}[t] & 1 + t\mathbb{C}[t] \end{pmatrix} \mid \det(g) = 1 \right\}$$

(respectively, corresponding unipotent cells) of $\mathrm{SL}_2(\mathcal{L})$. Hence we deal with $\mathrm{SL}_2(\mathcal{L})$ instead of $\widehat{\mathrm{SL}}_2(\mathcal{L})$. The torus H is the subgroup of $\mathrm{SL}_2(\mathcal{L})$ where the elements are the diagonal matrices of the form $\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}$. The Weyl group $W = \mathrm{Norm}(H)/H$ is generated by the two non-commuting involutions s_0 and s_1 .

Each $\varphi(P_{i,k})$ for $i = 0, 1$ and $k > 0$ is a regular function by computing explicit determinantal formulas. For $g = \begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix} \in U$, let T_g be the $\mathbb{Z} \times \mathbb{Z}$ matrix whose (M, N) entry is given by the residue formula

$$(T_g)_{M,N} = \mathrm{Res} \frac{g_{rs}}{t^{n-m+1}}$$

with $N = 2n + r$ and $M = 2m + s$ and with $n, m \in \mathbb{Z}$ and $r, s \in \{1, 2\}$. Let $\Delta_{k;i}^\sigma(g)$ be the determinant of the $k \times k$ submatrix of T_g whose row and column sets are given respectively by

$$\begin{aligned} \text{rows} &= \{k-i, k-i-2, k-i-4, k-i-6, \dots, 2-k-i\}, \\ \text{columns} &= \{k-i+1, k-i, k-i-1, k-i-2, \dots, 2-i\}. \end{aligned}$$

The following theorem is a special case of [Sc, Theorem 2].

Theorem III.3.3. $\varphi(P_{0,k}) = \Delta_{k;1}^\sigma$ and $\varphi(P_{1,k}) = \Delta_{k;0}^\sigma$ for any $k > 0$.

In the present case of the Weyl group elements $w_3 = s_0 s_1 s_0$ and $w_4 = s_0 s_1 s_0 s_1$, the corresponding affine unipotent varieties U^w are given by:

$$(7) \quad \begin{aligned} U^{w_3} &= \left\{ \left(\begin{array}{cc} 1+At & B \\ Dt+Et^2 & 1+ Ft \end{array} \right) \mid \begin{array}{l} A+F=BD \\ AF=BE \end{array} \quad E \neq 0 \right\} \\ U^{w_4} &= \left\{ \left(\begin{array}{cc} 1+At & B+Ct \\ Dt+Et^2 & 1+ Ft+Gt^2 \end{array} \right) \mid \begin{array}{l} A+F=BD \\ AF-CD=BE-G \end{array} \quad \begin{array}{l} AG=CE \\ G \neq 0 \end{array} \right\} \end{aligned}$$

In terms of the complex parameters $A, \dots, G$ we have

$$(8) \quad \begin{aligned} \varphi(P_{0,1}) &= \Delta_{1;1}^\sigma = D & \varphi(P_{1,2}) &= \Delta_{2;0}^\sigma = DF - E \\ \varphi(P_{0,3}) &= \Delta_{3;1}^\sigma = DEF - D^2G - E^2 & \varphi(P_{1,4}) &= \Delta_{4;0}^\sigma = G(DEF - D^2G - E^2) \end{aligned}$$

We are now ready to prove the crucial result on transcendence bases.

Proposition III.3.4. *The collections $\{\Delta_{1;1}^\sigma, \Delta_{2;0}^\sigma, \Delta_{3;1}^\sigma\}$ and $\{\Delta_{1;1}^\sigma, \Delta_{2;0}^\sigma, \Delta_{3;1}^\sigma, \Delta_{4;0}^\sigma\}$ are respectively transcendence bases for the rational function fields $\mathbb{C}(U^{w_3})$ and $\mathbb{C}(U^{w_4})$.*

Proof. Consider first the case of $\{\Delta_{1;1}^\sigma, \Delta_{2;0}^\sigma, \Delta_{3;1}^\sigma, \Delta_{4;0}^\sigma\}$ within $\mathbb{C}(U^{w_4})$. The transcendence degree of $\mathbb{C}(U^{w_4})$ over $\mathbb{C}$ is 4 since $\dim_{\mathbb{C}} U^{w_4}$ is 4. In this case any 4 rational functions which generate the field $\mathbb{C}(U^{w_4})$ must constitute a transcendence basis for $\mathbb{C}(U^{w_4})$. The global coordinates $A, B, C, D, E, F, G^{\pm 1}$ clearly generate $\mathbb{C}(U^{w_4})$, and we will be finished if we can express each global coordinate as a rational function in $\Delta_{1;1}^\sigma, \Delta_{2;0}^\sigma, \Delta_{3;1}^\sigma$, and $\Delta_{4;0}^\sigma$. To do this we introduce five auxiliary functions $\tilde{\Delta}_{1;1}^\sigma, \tilde{\Delta}_{2;0}^\sigma, \Psi, \Omega$, and Σ defined implicitly by the formulas

$$(9) \quad \begin{aligned} \tilde{\Delta}_{1;1}^\sigma \Delta_{1;1}^\sigma &= (\Delta_{2;0}^\sigma)^2 + \Delta_{3;1}^\sigma & \tilde{\Delta}_{2;0}^\sigma \Delta_{2;0}^\sigma &= (\Delta_{1;1}^\sigma)^2 \Delta_{4;0}^\sigma + (\Delta_{3;1}^\sigma)^2 \\ \Psi \Delta_{2;0}^\sigma &= (\tilde{\Delta}_{1;1}^\sigma)^2 + \Delta_{4;0}^\sigma & \Omega \Delta_{1;1}^\sigma &= (\tilde{\Delta}_{2;0}^\sigma)^2 + (\Delta_{3;1}^\sigma)^3 \\ \Sigma \tilde{\Delta}_{2;0}^\sigma &= (\Delta_{3;1}^\sigma)^4 \Delta_{4;0}^\sigma + \Omega^2 \end{aligned}$$

These are exactly the cluster variables obtained from the initial cluster from at most two mutations. Evidently these five functions can be rationally expressed in terms of $\Delta_{1;1}^\sigma, \Delta_{2;0}^\sigma, \Delta_{3;1}^\sigma$, and $\Delta_{4;0}^\sigma$. Moreover, after using the equations (8) and carrying out the divisions arising in solving the above system, each function can be written as a polynomial in the global coordinates $A, \dots, G^{\pm 1}$:

$$(10) \quad \begin{aligned} \tilde{\Delta}_{1;1}^\sigma &= DF^2 - EF - DG \\ \tilde{\Delta}_{2;0}^\sigma &= E(DEF - D^2G - E^2) \\ \Psi &= (F^2 - G)(DF - E) - DFG \\ \Omega &= (EF - DG)(DEF - D^2G - E^2)^2 \\ \Sigma &= (EF^2 - DFG - EG)(DEF - D^2G - E^2)^3 \end{aligned}$$

The following identities can be easily checked by using the relations (7).

$$(11) \quad \begin{aligned} A &= \frac{\Omega \tilde{\Delta}_{2;0}^\sigma}{\left(\Delta_{3;1}^\sigma\right)^4} & B &= \frac{\Sigma}{\left(\Delta_{3;1}^\sigma\right)^4} & C &= \frac{\Omega \Delta_{4;0}^\sigma}{\left(\Delta_{3;1}^\sigma\right)^4} & D &= \Delta_{1;1}^\sigma \\ E &= \frac{\tilde{\Delta}_{2;0}^\sigma}{\Delta_{3;1}^\sigma} & F &= \frac{\Sigma \Delta_{1;1}^\sigma + \Omega \tilde{\Delta}_{2;0}^\sigma}{\left(\Delta_{3;1}^\sigma\right)^4} & G^{\pm 1} &= \left(\frac{\Delta_{4;0}^\sigma}{\Delta_{3;1}^\sigma}\right)^{\pm 1} \end{aligned}$$

Thus $\{\Delta_{1;1}^\sigma, \Delta_{2;0}^\sigma, \Delta_{3;1}^\sigma, \Delta_{4;0}^\sigma\}$ is a transcendence basis.

The case of $\{\Delta_{1;1}^\sigma, \Delta_{2;0}^\sigma, \Delta_{3;1}^\sigma\}$ is handled similarly. Over the unipotent cell U^{w_3} the global coordinates $A, B, D, E^{\pm 1}, F$ are rationally expressed as

$$A = \frac{\tilde{\Delta}_{1;1}^\sigma}{\left(\Delta_{2;0}^\sigma\right)^2} \quad B = \frac{\left(\tilde{\Delta}_{1;1}^\sigma\right)^2}{\left(\Delta_{2;0}^\sigma\right)^3} \quad D = \Delta_{1;1}^\sigma \quad E^{\pm 1} = \left(\frac{\Delta_{3;1}^\sigma}{\Delta_{2;0}^\sigma}\right)^{\pm 1} \quad F = \frac{\tilde{\Delta}_{1;1}^\sigma}{\Delta_{2;0}^\sigma}$$

from which it follows that $\{\Delta_{1;1}^\sigma, \Delta_{2;0}^\sigma, \Delta_{3;1}^\sigma\}$ generates the rational function field $\mathbb{C}(U^{w_3})$. The transcendence degree of $\mathbb{C}(U^{w_3})$ is 3 and consequently the collection $\{\Delta_{1;1}^\sigma, \Delta_{2;0}^\sigma, \Delta_{3;1}^\sigma\}$ is a transcendence basis. $\square$

Theorem III.3.5. *For $w = w_3 = s_0 s_1 s_0$ or $w = w_4 = s_0 s_1 s_0 s_1$, the coordinate ring $\mathbb{C}[U^w]$ is a cluster algebra modelled by a strong cluster map $\varphi_w: \text{Sub } \Lambda / I_w \rightarrow \mathbb{C}[U^w]$ for the standard component of the cluster tilting graph of $\text{Sub } \Lambda / I_w$ with all coefficients inverted.*

Proof. We treat the case $w = w_4$, and leave the other easier case to the reader. Since $\varphi_{w_4}: \text{Sub } \Lambda / I_{w_4} \rightarrow \mathbb{C}[U^{w_4}]$ is a strong cluster map by Proposition III.3.4, we have a cluster algebra $\mathbf{A}^{w_4} \subset \mathbb{C}[U^{w_4}]$ where we invert all coefficients.

We start with proving the inclusion $\mathbf{A}^{w_4} \subset \mathbb{C}[U^{w_4}]$. By construction the functions $\Delta_{1;1}^\sigma, \Delta_{2;0}^\sigma, \Delta_{3;1}^\sigma$, and $\Delta_{4;0}^\sigma$ are regular. The coefficients $\Delta_{3;1}^\sigma$ and $\Delta_{4;0}^\sigma$ are invertible, so we must verify that their inverses are in $\mathbb{C}[U^{w_4}]$. Put

$$\Delta_{3;1} = BCF - B^2G - C^2, \quad \Delta_{4;0} = G(BCF - B^2G - C^2).$$

Using relations (7) one has $\Delta_{3;1}^\sigma \Delta_{3;1} = \Delta_{4;0}^\sigma \Delta_{4;0} = G^4$. The function $G \neq 0$ over U^{w_4} , and consequently $\Delta_{3;1}^\sigma$ and $\Delta_{4;0}^\sigma$ are invertible with inverses given by

$$\left(\Delta_{3;1}^\sigma\right)^{-1} = \frac{\Delta_{3;1}}{G^4} \quad \left(\Delta_{4;0}^\sigma\right)^{-1} = \frac{\Delta_{4;0}}{G^4}.$$

The two exchange relations for the initial seed are precisely the first two relations given in (9), namely

$$\tilde{\Delta}_{1;1}^\sigma \Delta_{1;1}^\sigma = \left(\Delta_{2;0}^\sigma\right)^2 + \Delta_{3;1}^\sigma, \quad \tilde{\Delta}_{2;0}^\sigma \Delta_{2;0}^\sigma = \left(\Delta_{1;1}^\sigma\right)^2 \Delta_{4;0}^\sigma + \left(\Delta_{3;1}^\sigma\right)^2$$

and we have seen in (10) that we have the following expressions for the cluster variables $\tilde{\Delta}_{1;1}^\sigma$ and $\tilde{\Delta}_{2;0}^\sigma$:

$$(12) \quad \tilde{\Delta}_{1;1}^\sigma = DF^2 - EF - DG, \quad \tilde{\Delta}_{2;0}^\sigma = E(DEF - D^2G - E^2)$$

which are clearly regular functions on U^{w_4} . Define $x_1 = \Delta_{1;1}^\sigma$, $x_2 = \Delta_{2;0}^\sigma$, $x_3 = \Delta_{3;1}^\sigma$, $x_4 = \Delta_{4;0}^\sigma$, and define also $\tilde{x}_1 = \tilde{\Delta}_{1;1}^\sigma$, and $\tilde{x}_2 = \tilde{\Delta}_{2;0}^\sigma$. Select an arbitrary cluster variable x in $\mathbf{A}^{w_4}$. By the Laurent phenomenon we know that x can be expressed as a polynomial in $x_1^{\pm 1}, x_2^{\pm 1}, x_3$, and x_4 . From this it follows that x is regular on the Zariski open set $\mathcal{U}$ of U^{w_4} defined by

$$\mathcal{U} = \left\{ g \in U^{w_4} \mid x_i(g) \neq 0 \text{ for } i = 1, 2 \right\}$$

Again, by the Laurent phenomenon, x can also be expressed, simultaneously, as a polynomial in both $\{\tilde{x}_1^{\pm 1}, x_2^{\pm 1}, x_3, x_4\}$ and $\{x_1^{\pm 1}, \tilde{x}_2^{\pm 1}, x_3, x_4\}$. Consequently x is simultaneously regular on both of the Zariski open subsets

$$\begin{aligned}\mathcal{U}^{(1)} &= \left\{ g \in U^{w_4} \mid \tilde{x}_1(g) \neq 0 \text{ and } x_2(g) \neq 0 \right\} \\ \mathcal{U}^{(2)} &= \left\{ g \in U^{w_4} \mid x_1(g) \neq 0 \text{ and } \tilde{x}_2(g) \neq 0 \right\}\end{aligned}$$

Taken all together we conclude that x is regular over the union $\mathcal{U} \cup \mathcal{U}^{(1)} \cup \mathcal{U}^{(2)}$. The complement V of this union inside U^{w_4} consists of those points g for which $x_1(g) = \tilde{x}_1(g) = 0$ and/or $x_2(g) = \tilde{x}_2(g) = 0$. Using the expressions for $\tilde{x}_1$ and $\tilde{x}_2$ given in (8) we see that

$$V = \left\{ \begin{pmatrix} 1 & B \\ Et^2 & 1 + Gt^2 \end{pmatrix} \mid \begin{array}{l} BE = G \\ G \neq 0 \end{array} \right\} \cup \left\{ \begin{pmatrix} 1 & Ct \\ Dt & 1 + Gt^2 \end{pmatrix} \mid \begin{array}{l} CD = G \\ G \neq 0 \end{array} \right\}$$

The only hypothetical singularities of x must lie in V , and since the codimension of V inside U^{w_4} is clearly 2, it follows that x has no singularities and is thus regular on all of U^{w_4} . The cluster algebra $\mathbf{A}^{w_4}$ is generated by the coefficients and cluster variables and so we conclude that $\mathbf{A}^{w_4} \subset \mathbb{C}[U^{w_4}]$.

The reverse inclusion $\mathbb{C}[U^{w_4}] \subset \mathbf{A}^{w_4}$ follows from the fact the functions x , Ω , and Σ defined previously are also cluster variables of $\mathbf{A}^{w_4}$ and the fact that the global coordinates $A, \dots, G^{\pm 1}$ are generated inside of $\mathbf{A}^{w_4}$ by using the expressions in (11). $\square$

Finally we show the cluster graphs for $\mathbb{C}[U^{w_3}]$ and $\mathbb{C}[U^{w_4}]$. Then $\mathbb{C}[U^{w_3}]$ has a seed consisting of a cluster $\{\Delta_{1;1}^\sigma\}$, coefficients $\{\Delta_{2;0}^\sigma, \Delta_{3;1}^\sigma\}$ and the quiver

$$\begin{array}{ccc} & \Delta_{2;0}^\sigma & \\ \nearrow & & \nwarrow \\ \Delta_{1;1}^\sigma & \longleftarrow & \Delta_{3;1}^\sigma \end{array}$$

which is the quiver of $\text{End}_\Lambda(T_3)$, where we drop the arrows between coefficients. The cluster graph is

$$\{\Delta_{1;1}^\sigma\} - \{\tilde{\Delta}_{1;1}^\sigma\}$$

where the only other cluster variable $\tilde{\Delta}_{1;1}^\sigma$ is determined by $\Delta_{1;1}^\sigma \tilde{\Delta}_{1;1}^\sigma = (\Delta_{2;0}^\sigma)^2 + \Delta_{3;1}^\sigma$. The cluster type of $\mathbb{C}[U^{w_3}]$ is A_1 .

Similarly, $\mathbb{C}[U^{w_4}]$ has a seed consisting of a cluster $\{\Delta_{1;1}^\sigma, \Delta_{2;0}^\sigma\}$, coefficients $\{\Delta_{3;1}^\sigma, \Delta_{4;0}^\sigma\}$ and the quiver

$$\begin{array}{ccccc} & \Delta_{2;0}^\sigma & \longleftarrow & \Delta_{4;0}^\sigma & \\ \nearrow & & & & \nwarrow \\ \Delta_{1;1}^\sigma & \longleftarrow & \Delta_{3;1}^\sigma & & \end{array}$$

The cluster graph is

$$\dots - \dots - \{\Delta_{1;1}^\sigma, \Delta_{2;0}^\sigma\} - \dots - \dots$$

The cluster type of $\mathbb{C}[U^{w_4}]$ is $\widehat{A}_1$.

Note that this gives an example of a substructure of a cluster structure coming from the inclusion $\text{Sub } \Lambda/I_{w_3} \subset \text{Sub } \Lambda/I_{w_4}$, and a cluster map such that we get a subcluster algebra of a cluster algebra, namely $\mathbb{C}[U^{w_3}]$ as a subcluster algebra of $\mathbb{C}[U^{w_4}]$.

REFERENCES

- [AHK] Angeleri Hügel L., Happel D., Krause H. *Handbook of tilting theory*, London Mathematical Society Lecture Note Series 332. Cambridge: Cambridge University Press
- [ABS] Assem I., Brüstle T., Schiffler R. *Cluster-tilted algebras as trivial extensions*, preprint arxiv:math.RT/0601537
- [ASS] Assem I., Simson D., Skowroński A. *Elements of the representation theory of associative algebras. Vol. 1. Techniques of representation theory*, London Mathematical Society Student Texts, 65. Cambridge University Press, Cambridge, 2006
- [A1] Auslander M. *Coherent functors*, 1966 Proc. Conf. Categorical Algebra (La Jolla, Calif.), (1965), 189–231
- [A2] Auslander M. *Applications of morphisms determined by modules*, Representation theory of algebras (Proc. Conf., Temple Univ., Philadelphia, Pa., 1976), pp. 245–327. Lecture Notes in Pure Appl. Math., Vol. 37, Dekker, New York, 1978
- [AR1] Auslander M., Reiten I. *Applications of contravariantly finite subcategories*, Adv. Math. 86 (1991), no.1, 111–152
- [AR2] Auslander M., Reiten I. *DTr-periodic modules and functors*, Repr. theory of algebras (Cocoyoc 1994) CMS Conf. Proc., vol. 18, Amer. Math. Soc., Providence, RI (1996), 39–50
- [ARS] Auslander M., Reiten I., Smalø S. O. *Representation theory of Artin algebras*, Cambridge Studies in Advanced Mathematics, 36. Cambridge University Press, Cambridge, 1997
- [AS] Auslander M., Smalø S.O. *Preprojective modules over Artin algebras*, J. Algebra 66 (1980), no. 1, 61–122
- [BGL] Baer D., Geigle W., Lenzing H. *The preprojective algebra of a tame hereditary Artin algebra*, Comm. Algebra 15 (1987), no. 1-2, 425–457
- [BFZ] Berenstein A., Fomin S., Zelevinsky A. *Cluster algebras. III. Upper bounds and double Bruhat cells*, Duke Math. J. 126 (2005), no. 1, 1–52
- [BZ] Berenstein A., Zelevinsky A. *Total positivity in Schubert varieties*, Comment. Math. Helv. 72 (1997), no. 1, 128–166
- [BL] Billey S., Lakshmibai V. *Singular loci of Schubert varieties*, Progress in Mathematics, 182. Birkhäuser Boston, Inc., Boston, MA, 2000
- [BB] Björner A., Brenti F. *Combinatorics of Coxeter groups*. Graduate Texts in Maths. 231. Springer, New York, 2005
- [B] Bocklandt R. *Graded Calabi Yau Algebras of dimension 3*, preprint arxiv:math.RA/0603558
- [BBK] Brenner S., Butler M. and King A., *Periodic algebras which are almost Koszul*, Algebr. Represent. Theory 5 (2002), no. 4, 331–367
- [BM] Buan A., Marsh R. *Cluster-tilting theory*, In Trends in Representation Theory of Algebras and Related Topics, Workshop August 11-14, 2004, Queretaro, Mexico. Contemporary Mathematics 406 (2006), 1–30
- [BMR1] Buan A., Marsh R., Reiten I. *Cluster-tilted algebras*, Trans. Amer. Math. Soc. 359 (2007), no. 1, 323–332
- [BMR2] Buan A., Marsh R., Reiten I. *Cluster mutation via quiver representations*, to appear in Comm. Math. Helv. arxiv:math.RT/0412077
- [BMRT] Buan A., Marsh R., Reiten I., Todorov G. *Clusters and seeds for acyclic cluster algebras* with an appendix by Buan A., Caldero P., Keller B., Marsh R., Reiten I., Todorov G., Proc. Amer. Math. Soc. 135, No. 10 (2007), 3049–3060
- [BMRRT] Buan A., Marsh R., Reineke M., Reiten I., Todorov G. *Tilting theory and cluster combinatorics*, Adv. Math. 204 (2006), no. 2, 572–618
- [BR] Buan A., Reiten I. *Acyclic quivers of finite mutation type*, International Mathematics Research Notices, Article ID 12804 (2006), 1–10
- [BIKR] Burban I., Iyama O., Keller B., Reiten I. *Cluster tilting for one-dimensional hypersurface singularities*, to appear in Adv. Math., arxiv:0704.1249
- [CC] Caldero P., Chapoton F. *Cluster algebras as Hall algebras of quiver representations*, Comment. Math. Helv. 81 (2006), no. 3, 595–616
- [CCS] Caldero P., Chapoton F., Schiffler R. *Quivers with relations arising from clusters (A_n case)*, Trans. Amer. Math. Soc. 358 (2006), no. 3
- [CK1] Caldero P., Keller B. *From triangulated categories to cluster algebras*, to appear in Invent. Math., arxiv: math.RT/0506018
- [CK2] Caldero P., Keller B. *From triangulated categories to cluster algebras II*, to appear in Annales Scientifiques de l'Ecole Normale Supérieure, arxiv: math.RT/0510251
- [CB] Crawley-Boevey W. *On the exceptional fibres of Kleinian singularities*, Amer. J. Math. 122 (2000), 1027–1037
- [EH] Erdmann K., Holm T. *Maximal n -orthogonal modules for selfinjective algebras*, to appear in Proc. Amer. Math. Soc., arXiv:math.RT/0603672

- [FZ1] Fomin S., Zelevinsky A., *Double Bruhat cells and total positivity*, J. Amer. Math. Soc. 12, (1999), 335–380
- [FZ2] Fomin S., Zelevinsky A. *Cluster Algebras I: Foundations*, J. Amer. Math. Soc. 15, no. 2 (2002), 497–529
- [FZ3] Fomin S., Zelevinsky A. *Cluster algebras II: Finite type classification*, Invent. Math. **154**, no. 1 (2003), 63–121
- [FZ4] Fomin S., Zelevinsky A. *Cluster algebras IV: Coefficients*, Compositio Mathematica 143, no. 1 (2007), 112–164
- [CE] Cartan H., Eilenberg S. *Homological algebra*, Princeton University Press, Princeton, N. J., 1956
- [GLS1] Geiss C., Leclerc B., Schröer J. *Rigid modules over preprojective algebras*, Invent. Math. 165 (2006), no. 3, 589–632
- [GLS2] Geiss C., Leclerc B., Schröer J. *Semicanonical bases and preprojective algebras*, Ann. Sci. Ecole Norm. Sup. (4) 38 (2005), no. 2, 193–253
- [GLS3] Geiss C., Leclerc B., Schröer J. *Auslander algebras and initial seeds for cluster algebras*, to appear in J. London Math. Soc., arxiv:math.RT/0506405
- [GLS4] Geiss C., Leclerc B., Schröer J. *Semicanonical bases and preprojective algebras II: A multiplication formula*, to appear in Compositio Mathematica, arxiv:math.RT/0509483
- [GLS5] Geiss C., Leclerc B., Schröer J. *Partial flag varieties and preprojective algebras*, preprint arxiv: math.RT/0609138
- [GLS6] Geiss C., Leclerc B., Schröer J. *Rigid modules over preprojective algebras II: The Kac-Moody case*, preprint arxiv: math.RT/0703039
- [H1] Happel D. *Triangulated categories in the representation theory of finite-dimensional algebras*, London Mathematical Society Lecture Note Series, 119. Cambridge University Press, Cambridge, 1988
- [H2] Happel D. *On Gorenstein algebras*, Representation theory of finite groups and finite-dimensional algebras (Bielefeld, 1991), 389–404, Progr. Math., 95, Birkhäuser, Basel, 1991
- [HU] Happel D., Unger L. *On a partial order of tilting modules*, Algebr. Represent. Theory 8 (2005), no. 2, 147–156
- [Hu] Hubery A. *Acyclic cluster algebras via Ringel-Hall algebras*, manuscript
- [Ig] Igusa K. *Notes on the no loops conjecture*, J. Pure Appl. Algebra 69 (1990), no. 2, 161–176
- [I1] Iyama O. *Higher-dimensional Auslander-Reiten theory on maximal orthogonal subcategories*, Adv. Math. 210 (1), (2007), 22–50
- [I2] Iyama O. *Auslander correspondence*, Adv. Math. 210 (1), (2007), 51–82
- [I3] Iyama O. *d-Calabi-Yau algebras and d-cluster tilting subcategories*, work in progress
- [IR] Iyama O., Reiten I. *Fomin-Zelevinsky mutation and tilting modules over Calabi-Yau algebras*, to appear in Amer. J. Math., arxiv:math.RT/0605136
- [IY] Iyama O., Yoshino Y. *Mutation in triangulated categories and rigid Cohen-Macaulay modules*, preprint arxiv:math.RT/0607736
- [IT] Ingalls C., Thomas H. *Noncrossing partitions and representations of quivers*, preprint arxiv:math.RT/0612219
- [KP1] Kac V.G., Peterson D.H. *Defining relations of certain infinite-dimensional groups*, Proceedings of the Cartan conference, Lyon 1984, Asterisque (1985), Numero hors serie, 165–208
- [KP2] Kac V.G., Peterson D.H. *Regular functions on certain infinite dimensional groups*, Progress in Math. 36 (1983), 141–166
- [Kel1] Keller B. *On triangulated orbit categories*, Documenta Math. 10 (2005), 551–581
- [Ke2] Keller B. *Derived categories and their uses*, Handbook of algebra, Vol. 1, 671–701, North-Holland, Amsterdam, 1996
- [Ke3] Keller B. *Chain complexes and stable categories*, Manuscripta Math. 67, (1990), no. 4, 379–417
- [KR1] Keller B., Reiten I. *Cluster-tilted algebras are Gorenstein and stably Calabi-Yau*, Adv. Math. 211 (1), (2007), 123–151
- [KR2] Keller B., Reiten I. *Acyclic Calabi-Yau categories*, preprint arxiv:math.RT/0610594
- [KZ] König S., Zhu B. *From triangulated categories to abelian categories - cluster tilting in a general framework*, preprint arxiv:math.RT/0605100
- [Le] Lenzing H. *Nilpotente Elemente in Ringen von endlicher globaler Dimension*, Math. Z. 108, 1969, 313–324
- [Lul1] Lusztig G. *Quivers, perverse sheaves, and quantized enveloping algebras*, J. Amer. Math. Soc. 4 (1991), no. 2, 365–421
- [Lu2] Lusztig G. *Semicanonical bases arising from enveloping algebras*, Adv. Math. 151 (2000), no. 2, 129–139
- [MRZ] Marsh, R., Reineke, M., Zelevinsky, A. *Generalized associahedra via quiver representations*, Trans. Amer. Math. Soc. 355 (2003), no. 10, 4171–4186
- [P] Palu Y. *Cluster characters for triangulated 2-Calabi-Yau categories*, preprint arxiv:math.RT/0703540

- [PS] Pressley A., Segal G. *Loop groups*, Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1986
- [RV] Reiten I., Van den Bergh M. *Noetherian hereditary abelian categories satisfying Serre duality*, J. Amer. Math. Soc. 15 (2002), no. 2, 295–366
- [Ri] Rickard J. *Morita theory for derived categories*, J. London Math. Soc. (2), 29 (1989), no.3, 436–456
- [Rin1] Ringel C.M. *Tame algebras and integral quadratic forms*, Lecture Notes in Mathematics, 1099. Springer-Verlag, Berlin, 1984
- [Rin2] Ringel C.M. *Some Remarks Concerning Tilting Modules and Tilted Algebras. Origin. Relevance. Future. (An appendix to the Handbook of Tilting Theory)*, Cambridge University Press (2007), LMS Lecture Notes Series 332
- [Sc] Scott J. *Block-Toeplitz determinants, chess tableaux, and the type $\hat{A}_1$ Geiss-Leclerc-Schröer φ -map*, preprint arXiv:0707.3046
- [Sm] Smalö S. O. *Torsion theories and tilting modules*, Bull. London Math. Soc. 16 (1984), no. 5, 518–522
- [T] Tabuada G. *On the structure of Calabi-Yau categories with a cluster tilting subcategory*, Doc. Math. 12 (2007), 193–213
- [Ye] Yekutieli A. *Dualizing complexes, Morita equivalence and the derived Picard group of a ring*, J. London Math. Soc. (2) 60 (1999), no. 3, 723–746
- [Yo] Yoshino Y. *Cohen-Macaulay modules over Cohen-Macaulay rings*, London Mathematical Society Lecture Note Series, 146. Cambridge University Press, Cambridge, 1990

INSTITUTT FOR MATEMATISKE FAG, NORGES TEKNISK-NATURVITENSKAPELIGE UNIVERSITET, N-7491 TRONDHEIM, NORWAY
E-mail address: `aslakb@math.ntnu.no`

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, CHIKUSA-KU, NAGOYA, 464-8602 NAGOYA, JAPAN
E-mail address: `iyama@math.nagoya-u.ac.jp`

INSTITUTT FOR MATEMATISKE FAG, NORGES TEKNISK-NATURVITENSKAPELIGE UNIVERSITET, N-7491 TRONDHEIM, NORWAY
E-mail address: `idunr@math.ntnu.no`

DEPT. OF PURE MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT, UNITED KINGDOM, ENGLAND
E-mail address: `jscott@maths.leeds.ac.uk`